



National
Comprehensive
Cancer
Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Breast Cancer Screening and Diagnosis

Version 1.2017 — June 2, 2017

NCCN.org

Continue



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 1.2017 Panel Members

Breast Cancer Screening and Diagnosis

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

***Therese B. Bevers, MD/Chair** **Ⓟ**
The University of Texas
MD Anderson Cancer Center

***Mark Helvie, MD/Vice-Chair** **Ⓟ** **Ⓢ**
University of Michigan
Comprehensive Cancer Center

Ermelinda Bonaccio, MD **Ⓟ**
Roswell Park Cancer Institute

Kristine E. Calhoun, MD **Ⓢ**
University of Washington/
Seattle Cancer Care Alliance

Mary B. Daly, MD, PhD **†**
Fox Chase Cancer Center

William B. Farrar, MD **Ⓢ**
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Judy E. Garber, MD, MPH **†**
Dana-Farber/Brigham and
Women's Cancer Center

Richard Gray, MD **Ⓢ**
Mayo Clinic Cancer Center

Caprice C. Greenberg, MD, MPH **Ⓢ**
University of Wisconsin
Carbone Cancer Center

Rachel Greenup, MD, MPH **Ⓢ**
Duke Cancer Institute

Nora M. Hansen, MD **Ⓢ**
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

NCCN
Mary Anne Bergman
Rashmi Kumar, PhD

Randall E. Harris, MD, PhD **Ⓟ** **Ⓢ**
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Alexandra S. Heerdt, MD **Ⓢ**
Memorial Sloan Kettering Cancer Center

Teresa Helsten, MD **†**
UC San Diego Moores Cancer Center

Linda Hodgkiss, MD **Ⓟ**
St. Jude Children's Research Hospital/
The University of Tennessee Health
Science Center

Tamarya L. Hoyt, MD **Ⓟ**
Vanderbilt-Ingram Cancer Center

John G. Huff, MD **Ⓟ**
Vanderbilt-Ingram Cancer Center

Debra M. Ikeda, MD **Ⓟ**
Stanford Cancer Institute

Lisa Jacobs, MD **Ⓢ**
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Constance Dobbins Lehman, MD, PhD
Massachusetts General Hospital
Cancer Center

Barbara Monsees, MD **Ⓟ**
Siteman Cancer Center at Barnes-
Jewish Hospital and Washington
University School of Medicine

Bethany L. Niell, MD, PhD **Ⓟ**
Moffitt Cancer Center

Catherine C. Parker, MD **Ⓢ**
University of Alabama at Birmingham
Comprehensive Cancer Center

Mark Pearlman, MD **Ⓢ** **Ⓢ**
University of Michigan
Comprehensive Cancer Center

Liane Philpotts, MD **Ⓟ**
Yale Cancer Center/Smilow Cancer Hospital

Laura B. Shepardson, MD **Ⓟ**
Case Comprehensive Cancer Center/
University Hospitals Seidman Cancer Center
and Cleveland Clinic Taussig Cancer Institute

Mary Lou Smith, JD, MBA **Ⓢ**
Research Advocacy Network

Matthew Stein, MD **Ⓟ**
Huntsman Cancer Institute
at the University of Utah

Lusine Tumyan, MD **Ⓟ**
City of Hope Comprehensive Cancer Center

Cheryl Williams, MD **Ⓟ**
Fred & Pamela Buffett Cancer Center

[NCCN Guidelines Panel Disclosures](#)

§ Radiation oncology/Radiotherapy
Ⓢ Surgery/Surgical oncology
† Medical oncology
‡ Hematology/Hematology oncology
Ⓟ Internist/Internal medicine, including family
practice, preventive management
Ⓢ Gynecologic oncology/Gynecology
Ⓟ Diagnostic/Interventional radiology
Ⓢ Pathology
Ⓢ Patient advocacy
* Discussion Section Writing Committee

Continue



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 1.2017 Table of Contents

Breast Cancer Screening and Diagnosis

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

[NCCN Breast Cancer Screening and Diagnosis Panel Members](#)

[Summary of the Guidelines Updates](#)

[Clinical Encounter Including Risk Assessment \(BSCR-1\)](#)

[Average Risk, Screening/Follow-Up \(BSCR-1\)](#)

[Increased Risk, Screening/Follow-Up \(BSCR-2\)](#)

[Symptomatic During Clinical Encounter, Presenting Signs/Symptoms \(BSCR-4\)](#)

- [Palpable Mass, Age ≥30 Years \(BSCR-5\)](#)
- [Palpable Mass, Age <30 Years \(BSCR-11\)](#)
- [Nipple Discharge, No Palpable Mass \(BSCR-13\)](#)
- [Asymmetric Thickening/Nodularity \(BSCR-14\)](#)
- [Skin Changes \(BSCR-15\)](#)

[Persistent or Severe Breast Pain \(BSCR-16\)](#)

[Mammographic Evaluation \(BSCR-19\)](#)

[Breast Screening Considerations \(BSCR-A\)](#)

[Risk Factors Used in the Modified Gail Model, Age ≥35 Years \(BSCR-B\)](#)

[Assessment Category Definitions \(BSCR-C\)](#)

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN Member Institutions, [click here: nccn.org/clinical_trials/physician.html](#).

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise specified.

See [NCCN Categories of Evidence and Consensus](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2017.



NCCN Guidelines Version 1.2017 Updates

Breast Cancer Screening and Diagnosis

Updates in Version 1.2017 of the NCCN Guidelines for Breast Cancer Screening from Version 2.2016 include:

GLOBAL: Footnotes changed throughout the algorithm.

BSCR-1

- "**Clinical encounter including risk assessment**" replaced "History and physical examination" as the first node on the page.
- "Negative physical exam" has been deleted from "Asymptomatic"
- "Positive physical exam" has been deleted from "Symptomatic"
- "Assess risk" has been included in the first node: *Clinical encounter including "risk assessment"*

Increased risk:

- 3rd bullet modified: ~~"Prior thoracic RT for Patients who receive thoracic RT younger than under 30 y (eg, mantle irradiation)"~~
- 5th bullet modified: "Women who have a lifetime risk $\geq 20\%$ based on history of LCIS or ADH/ALH" (Also for BSCR-3)

Footnotes

- "b" is new to the page: ~~"The FDA Medicare and insurers allow the patient direct access to scheduling for mammography."~~
- Modified footnote "c": "At minimum medical and family history should be obtained and clinical encounter should encompass ongoing risk assessment, risk reduction counseling, as well as clinical breast exam."
- "I" is new to the page: "Tomosynthesis improves cancer detection and reduces recall rates."

BSCR-2

- Modified: ~~"Prior thoracic Patient who receives thoracic RT between the ages for 10 and 30 y"~~

Screening/Follow-up

- 1st sub-bullet modified: "to begin at the age when identified as being at increased risk"

BSCR-3

Screening/Follow-up

- 1st and 2nd sub-bullets modified: "to begin at the age when identified as being at increased risk by Gail Model"

BSCR-4

- "Physical examination" deleted.
- Modified: "Symptomatic during clinical encounter ~~or positive findings on physical exam~~"

BSCR-5

Ultrasound Findings

- Top pathway: **Solid mass, Complex cystic and Solid mass and Complicated cyst** are new to the page.
- Modified: No ~~ultrasonographic~~ **ultrasonographic imaging** abnormality BI-RADS category 1.
- Bottom pathway: ~~Tissue Core needle~~ biopsy goes to BSCR-8, ~~BSCR-20~~ has been deleted.

Footnotes

- "p" modified: "Assess geographic correlation between clinical and imaging findings. If there is a lack of correlation, return to *mammogram findings: negative, benign or probably benign* for further workup of palpable lesion. If imaging findings correlate with the palpable finding, *subsequent workup will answer the problem.*"

BSCR-6

Ultrasound Finding/Palpable Mass

- 1st column modified to include "**Complex cystic and solid mass**"
- 3rd column bottom branch modified to include Physical exam \pm "**+ imaging**"...
- Bottom branch off "Non-simple cyst" has changed to "**Complicated cyst**":
- "Complex BI-RADS category 4 and Tissue biopsy" have been deleted.
- "Tissue biopsy" has been replaced with "**Core needle**" throughout the guidelines

Footnotes

- New footnote corresponding to "complicated cyst" BI-RADS category 3: "*In the context of numerous simple cysts, a complicated cyst may be considered a benign finding.*"
- "**t**" **Core needle biopsy preferred; in some circumstances needle aspiration may be sufficient.** This footnote corresponds to Core needle biopsy.
- "**p**": ~~Tissue sampling may be appropriate if clinically suspicious, aids in management, or is strongly desired by patient~~
- "**r**": ~~A complex cyst has both cystic and solid components.~~
- "**s**": ~~FNA and core (needle or vacuum-assisted) biopsy are both valuable. FNA requires cytologic expertise. Surgical excision is appropriate if unable to perform core needle biopsy or if strongly desired by patient.~~ Removed throughout the guidelines.

BSCR-7

- Title of the page has been modified: **Ultrasound Imaging Findings/Palable Mass** (Also for BSCR-12)
- 1st column modified: For age ≥ 30 y No ~~ultrasonographic~~ **ultrasonographic imaging** abnormality BI-RADS category 1 (Also for BSCR-12)



NCCN Guidelines Version 1.2017 Updates

Breast Cancer Screening and Diagnosis

Updates in Version 1.2017 of the NCCN Guidelines for Breast Cancer Screening from Version 2.2016 include:

BSCR-15

Presenting Signs/Symptoms

- Modified: "Clinical suspicion of inflammatory breast cancer *includes but is not limited to:*"
 - Peau d'orange (pitted or dimpled appearance of skin)
 - Skin thickening
 - Edema
 - Erythema"
- <30 y to *Ultrasound ± diagnostic mammogram* and ≥30 y to *Diagnostic mammogram + ultrasound* is new to the page.

Diagnostic Follow-up

- Modified: "Core needle biopsy (preferred) ± punch biopsy ~~or Surgical excision~~"

Footnotes

- "dd" has been removed from "Skin changes" and added to "Clinical suspicion of inflammatory breast cancer" and includes a link to an article on inflammatory breast cancer.
- "gg" is a new footnote corresponding to "biopsy" for the lower branch off
BI-RADS 4-5: "*Inflammatory breast cancer is a clinical diagnosis and is not dependent on a positive punch biopsy*"

BSCR-16

- "*Adequate clinical breast exams include the following: upright and supine position during inspection, and palpation of all components of the breast, axilla, and clavicular lymph node basins. Time spent on the palpable portion of the exam is associated with increased detection of palpable abnormalities. Location and distance from nipple facilitate geographic correlation with imaging findings (See BSCR-1)*" is new footnote to the page.

BSCR-17

Footnotes

- The following corresponds to "mammogram ≥30 y": "*There are some circumstances with low clinical suspicion; ultrasound would be preferred and may suffice for women 30–39 years of age.*"
- "hh": "~~Consider ultrasound if mammogram results do not explain the pain.~~"
- "ii": "~~Consider mammogram if ultrasound does not explain the pain.~~"

BSCR-18

Follow-up Evaluation

- Modified for BI-RADS category 1: "*Symptomatic management*" directs the reader to "*Follow-up screening*" ~~Clinical encounter every 6–12 months for 1–2 y (treatment if desired)~~
- Modified for BI-RADS category 3: "Follow-up with ~~age appropriate~~ diagnostic mammogram ± and/or ultrasound, 6–12 months for 1–2 y"

BSCR-19

Follow-up

Modified 5th node: "*After complete imaging evaluation tissue sampling by ultrasound-guided*" core needle biopsy

BSCR-20

- Page deleted, refer to BSCR-8.



NCCN Guidelines Version 1.2017 Updates

Breast Cancer Screening and Diagnosis

Updates in Version 1.2017 of the NCCN Guidelines for Breast Cancer Screening from Version 2.2016 include:

[BSCR-A 1 of 2](#)

- 1st bullet modified to include: "*Shared decision-making is encouraged based on a woman's values and preferences. (See Discussion)*"
- Former bullets 5 and 6 have been deleted: "~~but there is insufficient evidence to support routine supplemental screening in women with dense breasts and no other risk factors. Important outcomes are not yet established for supplemental screening; some states have passed legislation mandating patient notification of breast density:~~" and "~~There are several studies supporting the use of supplemental screening for breast cancer as an adjunct to screening mammography for women with dense breast tissue. Different modalities may be considered based on risk and patient values/preference.~~"
- 5th bullet is new to the page: "*For women with heterogeneous dense breasts and dense breast tissue, recommend counseling on the risks and benefits of supplemental screening.*"
- 7th bullet modified: "*Full-field digital mammography appears to benefit young women and women with dense breasts.*"
- 8th bullet modified: "~~Multiple studies show a combined use of digital mammography and tomosynthesis appears to improve cancer detection and decrease call back rates. Of note, most studies used double the dose of radiation. The radiation dose can be minimized by using synthetic 2-D reconstruction.~~"

[BSCR-A 2 of 2](#)

- 1st bullet modified: "*BRCA mutation, commence at age 25–29 y*"
- 2nd bullet modified: "*First-degree relative of BRCA carrier, but untested: commence at age 25–29 y*"
- 7th bullet added: "*NF1 and NBN have been added to the list of genes for >20% risk of breast cancer based on gene and/or risk level--ATM, CDH1, CHEK2, NF1, NBN, PALB2, PTEN, STK11, TP53*"

Footnote:

- Corresponding to the title of the page, "*Women with a history of breast cancer with these risk factors should consider supplemental screening*"

[BSCR-B](#)

- "*Needle biopsy counts for number of biopsies in the Gail Model*" is a new footnote to the page.




NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

SCREENING OR SYMPTOM CATEGORY

SCREENING/FOLLOW-UP^a



^aSee [Breast Screening Considerations \(BSCR-A\)](#).

^bMedicare and insurers allow the patient direct access to scheduling for mammography.

^cAt minimum medical and family history should be obtained and clinical encounter should encompass ongoing risk assessment, risk reduction counseling, as well as a clinical breast exam.

^dRefer to the [NCCN Guidelines for Breast Cancer Risk Reduction](#) for a detailed qualitative and quantitative assessment.

^eSee [NCCN Guidelines for Breast Cancer - Surveillance Section](#).

^fRisk models that are largely dependent on family history (eg, Claus, BRCAPRO, BOADICEA, Tyrer-Cuzick). See [NCCN Guidelines for Breast Cancer Risk Reduction](#).

^gSee [Risk Factors Used in the Modified Gail Model, Age ≥35 Years \(BSCR-B\)](#).

^hThere is variation in recommendations for initiation of screening for different genetic syndromes. See [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast and Ovarian](#).

ⁱRandomized trials comparing clinical breast exam versus no screening have not been performed. Rationale for recommending clinical encounter is to maximize earliest detection of breast cancers.

^jWomen should be familiar with their breasts and promptly report changes to their health care provider.

^kSee Mammographic Evaluation ([BSCR-19](#)).

^lTomosynthesis improves cancer detection and reduces recall rates.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

SCREENING OR SYMPTOM CATEGORY

SCREENING/FOLLOW-UP

Increased Risk:

Prior history of breast cancer → [See NCCN Guidelines for Breast Cancer](#) - Surveillance Section

OR

Women who have a lifetime risk
>20% as defined by models that are
largely dependent on family history^f

- Clinical encounter^{a,c,i} every 6–12 mo
 - to begin when identified as being at increased risk
 - Referral to genetic counseling if not already done
- Annual screening^a mammogram^k
 - to begin 10 years prior to the youngest family member but not less than age 30 y
 - Consider tomosynthesis^{a,l}
- Recommend annual breast MRI^m
 - to begin 10 years prior to youngest family member but not less than age 25 y
- Consider risk reduction strategies ([See NCCN Guidelines for Breast Cancer Risk Reduction](#))
- Breast awareness^j

OR

Patient who
receives thoracic
RT between the
ages of 10 and
30 y

Current age <25 y →

- Annual clinical encounter^{a,c,i}
 - beginning 8–10 y after RT
- Breast awareness^j

Current age ≥25 y →

- Clinical encounter^{a,c,i} every 6–12 mo
 - Begin 8–10 y after RT
- Annual screening^a mammogram^k
 - Begin 8–10 y after RT but not prior to age 25 y
 - Consider tomosynthesis^{a,l}
- Recommend annual breast MRI^m
 - Begin 8–10 y after RT but not prior to age 25 y
- Breast awareness^j

^a[See Breast Screening Considerations \(BSCR-A\)](#)

^cAt minimum medical and family history should be obtained and clinical encounter should encompass ongoing risk assessment, risk reduction counseling, as well as a clinical breast exam.

^fRisk models that are largely dependent on family history (eg, Claus, BRCAPRO, BOADICEA, Tyrer-Cuzick). [See NCCN Guidelines for Breast Cancer Risk Reduction](#)

ⁱRandomized trials comparing clinical breast exam versus no screening have not been performed. Rationale for recommending clinical encounter is to maximize earliest detection of breast cancers.

^jWomen should be familiar with their breasts and promptly report changes to their health care provider.

^kSee Mammographic Evaluation ([BSCR-19](#)).

^lTomosynthesis improves cancer detection and reduces recall rates.

^mHigh-quality breast MRI limitations include having: a need for a dedicated breast coil, the ability to perform biopsy under MRI guidance, experienced radiologists in breast MRI, and regional availability. Breast MRI is performed preferably days 7–15 of menstrual cycle for premenopausal women. MRI should be integrated with other breast imaging modalities.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

SCREENING OR SYMPTOM CATEGORY

SCREENING/FOLLOW-UP

Increased Risk:

Women ≥35 y with 5-year Gail Model risk of invasive breast cancer ≥1.7%⁹



- Clinical encounter^{a,c,i} every 6–12 mo
 - to begin when identified as being at increased risk by Gail Model
- Annual screening^a mammogram^k
 - to begin when identified as being at increased risk by Gail Model
 - Consider tomosynthesis^{a,l}
- Consider risk reduction strategies ([See NCCN Guidelines for Breast Cancer Risk Reduction](#))
- Breast awareness^j

OR

Women who have a lifetime risk ≥20% based on history of LCIS or ADH/ALH



- Clinical encounter^{a,c,i} every 6–12 mo
 - to begin at diagnosis of LCIS or ADH/ALH
- Annual screening^a mammogram^k
 - to begin at diagnosis of LCIS or ADH/ALH but not less than age 30 y
 - Consider tomosynthesis^{a,l}
- Consider annual MRI
 - to begin at diagnosis of LCIS or ADH/ALH but not less than age 25 y (based on emerging evidence)
- Consider risk reduction strategies ([See NCCN Guidelines for Breast Cancer Risk Reduction](#))
- Breast awareness

^a[See Breast Screening Considerations \(BSCR-A\)](#)

^cAt minimum medical and family history should be obtained and clinical encounter should encompass ongoing risk assessment, risk reduction counseling, as well as a clinical breast exam.

⁹[See Risk Factors Used in the Modified Gail Model, Age ≥35 Years \(BSCR-B\)](#).

ⁱRandomized trials comparing clinical breast exam versus no screening have not been performed. Rationale for recommending clinical encounter is to maximize earliest detection of breast cancers.

^jWomen should be familiar with their breasts and promptly report changes to their health care provider.

^kSee Mammographic Evaluation ([BSCR-19](#)).

^lTomosynthesis improves cancer detection and reduces recall rates.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

PRESENTING SIGNS/SYMPTOMS



Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017


Breast Cancer Screening and Diagnosis

PRESENTING SIGNS/SYMPTOMS

DIAGNOSTIC EVALUATION

WORKUP

ULTRASOUND FINDINGS



ⁿSee Assessment Category Definitions (BSCR-C).

^oThere are some clinical circumstances such as mass with low clinical suspicion or suspected simple cyst in which ultrasound would be preferred and may suffice for women 30–39 years of age. See Discussion section.

^pAssess geographic correlation between clinical and imaging findings. If there is a lack of correlation, return to mammogram findings: negative, benign or probably benign for further workup of palpable lesion. If imaging findings correlate with the palpable finding, subsequent workup will answer the problem.

^qConcordance is needed between clinical exam and imaging results. Consider therapeutic aspiration for persistent clinical symptoms.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

ULTRASOUND FINDINGS/PALPABLE MASS



ⁿSee Assessment Category Definitions (BSCR-C).

^rIn the context of numerous simple cysts, a complicated cyst may be considered a benign finding.

^sRound or oval, circumscribed mass containing low-level echoes without vascular flow, fulfilling most but not all criteria for simple cyst.

^tCore needle biopsy preferred; in some circumstances needle aspiration may be sufficient.

^uThere may be variability on the follow-up interval based on the level of suspicion.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

ULTRASOUND IMAGING FINDINGS/PALPABLE MASS



ⁿSee Assessment Category Definitions (BSCR-C).

^qConcordance is needed between clinical exam and imaging results. Consider therapeutic aspiration for persistent clinical symptoms.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis



^uThere may be variability on the follow-up interval based on the level of suspicion.

^vSelect patients may be suitable for monitoring in lieu of surgical excision (eg, FEA, papillomas, fibroepithelial lesions, radial scars).

^wOther histologies that may require additional tissue: mucin-producing lesions, potential phyllodes tumor, papillary lesions, radial scar, or histologies of concern to pathologist.

^xMultifocal/extensive LCIS involving >4 terminal ductal lobular units on a core biopsy may be associated with increased risk for invasive cancer on surgical excision. (Rendi MH, Dintzis SM, Lehman CD, et al. Lobular in-situ neoplasia on breast core needle biopsy: imaging indication and pathologic extent can identify which patients require excisional biopsy. Ann Surg Oncol 2012;19:914-921. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21861212>).

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

FOLLOW-UP EVALUATION



Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

ASPIRATE FINDINGS/PALPABLE MASS

FOLLOW-UP EVALUATION



^yRoutine cytology is not recommended.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

PRESENTING SIGNS/SYMPTOMS

DIAGNOSTIC EVALUATION



ⁿSee Assessment Category Definitions (BSCR-C).

^qConcordance is needed between clinical exam and imaging results. Consider therapeutic aspiration for persistent clinical symptoms.

^zIf suspicious or highly suggestive of malignancy obtain mammogram.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

ULTRASOUND IMAGING FINDINGS/PALPABLE MASS

FOLLOW-UP EVALUATION


ⁿSee Assessment Category Definitions (BSCR-C).

^{aa}Assess geographic correlation between clinical and imaging findings. If there is a lack of correlation, return to BI-RADS category 1-2 for further workup of palpable lesion. If imaging findings correlate with the palpable finding, subsequent workup will answer the problem.

^uThere may be variability on the follow-up interval based on the level of suspicion.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

PRESENTING SIGNS/ SYMPTOMS

DIAGNOSTIC FOLLOW-UP



ⁿSee Assessment Category Definitions (BSCR-C).

^{bb}A list of drugs that can cause nipple discharge (not all-inclusive): psychoactive drugs, antihypertensive medications, opiates, oral contraceptives, and estrogen.

^{cc}If BI-RADS category 3 finding is unrelated to nipple discharge, manage mammographic finding by [BSCR-19](#).


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis


ⁿSee Assessment Category Definitions (BSCR-C).

^oThere are some clinical circumstances such as mass with low clinical suspicion or suspected simple cyst, in which ultrasound would be preferred and may suffice for women 30–39 years of age. See Discussion section.

^{aa}Assess geographic correlation between clinical and imaging findings. If there is a lack of correlation, return to BI-RADS category 1-2 for further workup of palpable lesion. If imaging findings correlate with the palpable finding, subsequent workup will answer the problem.

^uThere may be variability on the follow-up interval based on the level of suspicion.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

PRESENTING SIGNS/ SYMPTOMS



ⁿSee Assessment Category Definitions (BSCR-C).

^oThere are some clinical circumstances such as mass with low clinical suspicion or suspected simple cyst, in which ultrasound would be preferred and may suffice for women 30–39 years of age. [See Discussion](#) section.

^{dd}This may represent serious disease of the breast and needs evaluation. (Dawood S, Merajver SD, Viens P, et al. International expert panel on inflammatory breast cancer: consensus statement for standardized diagnosis and treatment. Ann Oncol 2011;22(3):515-523. Available at <https://www.ncbi.nlm.nih.gov/pubmed/20603440>).

^{ee}If clinically of low suspicion for breast cancer or high suspicion for infection, a short trial (7–10 days) of antibiotics for mastitis may be indicated.

^{ff}A benign skin punch biopsy in a patient with a clinical suspicion of inflammatory breast cancer does not rule out malignancy. Further evaluation is recommended.

^{gg}Inflammatory breast cancer is a clinical diagnosis and is not dependent on a positive punch biopsy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

PRESENTING SIGNS AND SYMPTOMS

FOLLOW-UP EVALUATION



^{hh}Defined as 4 to 6 weeks duration prior to that, symptomatic management.

ⁱⁱAdequate clinical breast exams include the following: upright and supine position during inspection, and palpation of all components of the breast, axilla, and clavicular lymph node basins. Time spent on the palpable portion of the exam is associated with increased detection of palpable abnormalities. Location and distance from nipple facilitate geographic correlation with imaging findings. ([See BSCR-1](#)).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

PRESENTING SIGNS AND SYMPTOMS

FOLLOW-UP EVALUATION



^{jj}Assuming breast imaging screening is current.

^{kk}There are some circumstances with low clinical suspicion; ultrasound would be preferred and may suffice for women 30–39 years of age.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis



ⁿ[See Assessment Category Definitions \(BSCR-C\).](#)

^{||}If complicated cyst, consider aspiration.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis



ⁿ[See Assessment Category Definitions \(BSCR-C\).](#)

^uThere may be variability on the follow-up interval based on the level of suspicion.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

BREAST SCREENING CONSIDERATIONS

- **Women should be counseled regarding potential benefits, risks, and limitations of breast screening. Shared decision-making is encouraged based on a woman's values and preferences ([See Discussion](#)).**
- **Adequate clinical breast exams include the following: upright and supine position during inspection, and palpation of all components of the breast, axilla, and clavicular lymph node basins. Time spent on the palpable portion of the exam is associated with increased detection of palpable abnormalities. Location and distance from nipple facilitate geographic correlation with imaging findings.**
- **Consider severe comorbid conditions limiting life expectancy (eg, ≤ 10 years) and whether therapeutic interventions are planned.**
- **Upper age limit for screening is not yet established.**
- **For women with heterogeneous dense breasts and dense breast tissue, recommend counseling on the risks and benefits of supplemental screening.¹**
- **Dense breasts limit the sensitivity of mammography. Dense breasts are associated with an increased risk for breast cancer.**
- **Full-field digital mammography appears to benefit young women and women with dense breasts.²**
- **Multiple studies show tomosynthesis appears to improve cancer detection and decrease call back rates. Of note, most studies used double the dose of radiation. The radiation dose can be minimized by using synthetic 2-D reconstruction.**
- **Current evidence does not support the routine use of breast scintigraphy (eg, sestamibi scan) as a screening procedure, but there is emerging evidence that breast scintigraphy may improve detection of early breast cancers among women with mammographically dense breasts.**
- **Current evidence does not support the routine use of thermography or ductal lavage as screening procedures.**
- **In high-risk settings based on current evidence and considering the FDA warning³ (gadolinium-based contrast agents) we continue to recommend annual MRI in these select populations.**

¹Berg WA, Blume JD, Cormack JB, et al. Combined screening with ultrasound and mammography vs. mammography alone in women at elevated risk of breast cancer. JAMA 2008;299(18):2151-2163.

²Pisano ED, Gatsonis C, Hendrick E et al for the Digital Mammographic Imaging Screening Trial (DMIST) Investigators. Diagnostic performance of digital versus film mammography for breast cancer screening. N Engl J Med 2005;353:1773-1783.

³FDA Drug Safety Communication: FDA evaluating the risk of brain deposits with repeated use of gadolinium-based contrast agents for magnetic resonance imaging (MRI) <http://www.fda.gov/Drugs/DrugSafety/ucm455386.htm>

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

BREAST SCREENING CONSIDERATIONS

RECOMMENDATIONS FOR BREAST MRI SCREENING AS AN ADJUNCT TO MAMMOGRAPHY^{4,5} (FOR AGE TO BEGIN SCREENING EXCEPT WHERE NOTED BELOW: [SEE BSCR-2](#))

Recommend Annual MRI Screening (Based on Evidence):

- *BRCA* mutation, commence at age 25–29 y
- First-degree relative of *BRCA* carrier, but untested: commence at age 25–29 y
- Lifetime risk 20% or greater, as defined by models that are largely dependent on family history

Recommend Annual MRI Screening (Based on Expert Consensus Opinion):

- Radiation to chest between age 10 and 30 years
- Li-Fraumeni syndrome and first-degree relatives
- Bannayan–Riley–Ruvalcaba syndromes and first-degree relatives
- >20% risk of breast cancer based on gene and/or risk level--*ATM*, *CDH1*, *CHEK2*, *NF1*, *NBN*, *PALB2*, *PTEN*, *STK11*, *TP53*

Consider MRI screening for LCIS and ALH/ADH based on emerging evidence if lifetime risk $\geq 20\%$

Insufficient Evidence to Recommend for or Against MRI Screening:

- Lifetime risk 15%–20%, as defined by models that are largely dependent on family history
- Heterogeneously or extremely dense breast on mammography
- Women with a personal history of breast cancer, including ductal carcinoma in situ (DCIS)

Recommend Against MRI Screening (Based on Expert Consensus Opinion):

- Women at <15% lifetime risk

⁴Adapted with permission from John Wiley and Sons. Copyright ©2007 American Cancer Society. Saslow D, Boetes C, Burke W, et al. American Cancer Society Guidelines for Breast Cancer Screening with MRI as an Adjunct to Mammography. CA: Cancer J Clin 2007;57:75-89.

⁵Women with a history of breast cancer with these risk factors should consider supplemental screening.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

RISK FACTORS USED IN THE MODIFIED GAIL MODEL, AGE ≥35 Years¹

- Current age
- Age at menarche
- Age at first live birth or nulliparity
- Number of first-degree relatives with breast cancer
- Number of previous benign breast biopsies²
- Atypical hyperplasia in a previous breast biopsy
- Race³

For calculation of risk, based on the modified Gail Model, see
<http://www.cancer.gov/bcrisktool/Default.aspx>

¹For detailed information, see <http://www.cancer.gov/bcrisktool/Default.aspx>.

²Needle biopsy counts for number of biopsies in the Gail Model.

³The current Gail Model may not accurately assess breast cancer risk in non-Caucasian, non-Asian, and non-African American women.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

MAMMOGRAPHIC ASSESSMENT CATEGORY DEFINITIONS^{1,2}

BI-RADS® - MAMMOGRAPHY FINDINGS

A. Assessment Is Incomplete:

Category 0: Incomplete - Need Additional Imaging Evaluation and/or Prior Mammograms for Comparison:

There is a finding for which additional evaluation is needed. This is almost always used in a screening situation. Under certain circumstances this assessment category may be used in a diagnostic mammography report, such as when ultrasound equipment or personnel are not immediately available, or when the patient is unable or unwilling to wait for completion of a full diagnostic examination. A recommendation for additional imaging evaluation includes the use of spot compression (with or without magnification), special mammographic views, and ultrasound. Category 0 should not be used for diagnostic breast imaging findings that warrant further evaluation with MRI. Rather, the interpreting physician should issue a final assessment in a report that is made before the MRI examination is performed. In most circumstances and when feasible, if a mammography examination is not assessed as negative or benign, the current examination should be compared with prior examination(s). The interpreting physician should use judgment on how vigorously to attempt obtaining prior examinations, given the likelihood of success of such an endeavor and the likelihood that comparison will affect the final assessment. In this context, it is important to note that comparison with previous examination(s) may be irrelevant when a finding is inherently suspicious for malignancy.

Category 0 should be used for prior image comparison only when such comparison is required to make a final assessment. When category 0 is used in the context of awaiting prior examinations for comparison, there should be in place a tracking procedure guaranteeing with 100% reliability that a final assessment will be made within 30 days (preferably sooner) even if prior examinations do not become available. Some mammography practices may reasonably choose never to use category 0 in the context of awaiting prior examinations simply because they do not have a 100% reliable tracking procedure. If a mammography examination is assessed as category 0 in the context of awaiting prior examinations and then the prior examinations do become available, an addendum to the initial mammography report should be issued, including a revised assessment. For auditing purposes, the revised assessment should replace the initial assessment.

¹Mammography results are mandated to be reported using Final Assessment categories (Quality Mammography Standards: Final Rule. Federal Register. 1997;62:55988).

²Terminology in this table is reflective of the American College of Radiology (ACR). ACR-BI-RADS®--5th Edition. ACR Breast Imaging Reporting and Data System, Breast Imaging Atlas; BI-RADS. Reston VA. American College of Radiology, 2014. For more information, see www.acr.org. Reprinted with permission from the American College of Radiology. No other representation of this document is authorized without express, written permission from the American College of Radiology.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

MAMMOGRAPHIC ASSESSMENT CATEGORY DEFINITIONS^{1,2}

BI-RADS® - MAMMOGRAPHY FINDINGS

B. Assessment Is Complete - Final Assessment Categories:

Category 1: Negative:

There is nothing to comment on. This is a normal examination.

Category 2: Benign:

Like Category 1, this is a "normal" assessment, but here, the interpreter chooses to describe a benign finding in the mammography report. Involuting, calcified fibroadenomas, skin calcifications, metallic foreign bodies (such as core biopsy and surgical clips), and fat-containing lesions (such as oil cysts, lipomas, galactoceles, and mixed-density hamartomas) all have characteristically benign appearances and may be described with confidence. The interpreter may also choose to describe intramammary lymph nodes, vascular calcifications, implants, or architectural distortion clearly related to prior surgery while still concluding that there is no mammographic evidence of malignancy. On the other hand, the interpreter may choose not to describe such findings, in which case the examination should be assessed as negative (category 1).

Note that both category 1 and category 2 assessments indicate that there is no mammographic evidence of malignancy. Both should be followed by the management recommendation for routine mammography screening. The difference is that category 2 should be used when describing one or more specific benign mammographic findings in the report, whereas category 1 should be used when no such findings are described (even if such findings are present).

¹Mammography results are mandated to be reported using Final Assessment categories (Quality Mammography Standards: Final Rule. Federal Register. 1997;62:55988).

²Terminology in this table is reflective of the American College of Radiology (ACR). ACR-BI-RADS®--5th Edition. ACR Breast Imaging Reporting and Data System, Breast Imaging Atlas; BI-RADS. Reston VA. American College of Radiology, 2014. For more information, see www.acr.org. Reprinted with permission from the American College of Radiology. No other representation of this document is authorized without express, written permission from the American College of Radiology.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

MAMMOGRAPHIC ASSESSMENT CATEGORY DEFINITIONS^{1,2}

BI-RADS® - MAMMOGRAPHY FINDINGS

Category 3: Probably Benign:

A finding assessed using this category should have a $\leq 2\%$ likelihood of malignancy, but greater than the essentially 0% likelihood of malignancy of a characteristically benign finding. A probably benign finding is not expected to change over the suggested period of imaging surveillance, but the interpreting physician prefers to establish stability of the finding before recommending management limited to routine mammography screening.

There are several prospective clinical studies demonstrating the safety and efficacy of periodic mammographic surveillance instead of biopsy for specific mammographic findings.

Three specific findings are validated as being probably benign (the noncalcified circumscribed solid mass, the focal asymmetry, and solitary group of punctate calcifications). All the previously cited studies emphasize the need to conduct a complete diagnostic imaging evaluation before making a probably benign (category 3) assessment; hence, it is recommended not to render such an assessment in interpreting a screening mammography examination. The practice of rendering category 3 assessments directly from screening examination also has been shown to result in adverse outcomes: 1) unnecessary follow-up of many lesions that could have been promptly assessed as benign; and 2) delayed diagnosis of a small number of cancers that otherwise may have been smaller in size and less likely to be advanced in stage. Also, all the previously cited studies exclude palpable lesions, so the use of a probably benign assessment for a palpable lesion is not supported by robust scientific data, although there are two single-institution studies that do report successful outcomes for palpable lesions. Finally, because evidence from previously cited studies indicates the need for biopsy rather than continued surveillance when a probably benign finding increases in size or extent, it is not prudent to render a category 3 assessment when a finding that otherwise meets “probably benign” imaging criteria is either new or has increased in size or extent.

While the vast majority of probably benign findings are managed with an initial short-interval follow-up (6-month) examination followed by additional examinations until long-term (2- or 3-year) stability is demonstrated, there may be occasions in which a biopsy is done instead (patient preference or overriding clinical concern).

¹Mammography results are mandated to be reported using Final Assessment categories (Quality Mammography Standards: Final Rule. Federal Register. 1997;62:55988).

²Terminology in this table is reflective of the American College of Radiology (ACR). ACR-BI-RADS®--5th Edition. ACR Breast Imaging Reporting and Data System, Breast Imaging Atlas; BI-RADS. Reston VA. American College of Radiology, 2014. For more information, see www.acr.org. Reprinted with permission from the American College of Radiology. No other representation of this document is authorized without express, written permission from the American College of Radiology.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)

BSCR-C
3 OF 9



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

MAMMOGRAPHIC ASSESSMENT CATEGORY DEFINITIONS^{1,2}

BI-RADS® - MAMMOGRAPHY FINDINGS

Category 4: Suspicious:

This category is reserved for findings that do not have the classic appearance of malignancy but are sufficiently suspicious to justify a recommendation for biopsy. The ceiling for category 3 assessment is a 2% likelihood of malignancy and the floor for category 5 assessment is 95%, so category 4 assessments cover the wide range of likelihood of malignancy in between. Thus, almost all recommendations of breast interventional procedures will come from assessments made using this category. By subdividing category 4³ into 4A, 4B, and 4C, as recommended in Guidance chapter and using the cut point indicated therein, it is hoped that patients and referring clinicians will more readily make informed decisions on the ultimate course of action.

Category 5: Highly Suggestive of Malignancy:

These assessments carry a very high probability (≥95%) of malignancy. This category initially was established to involve lesions for which 1-stage surgical treatment was considered without preliminary biopsy, in an era when preoperative wire localization was the primary breast interventional procedure. Nowadays, given the widespread acceptance of imaging-guided percutaneous biopsy, 1-stage surgery is rarely, if ever, performed. Rather, current oncologic management almost always involves tissue diagnosis of malignancy via percutaneous tissue sampling to facilitate treatment options, such as when sentinel node biopsy is included in surgical management or when neoadjuvant chemotherapy is administered prior to surgery. Therefore, the current rationale for using a category 5 assessment is to identify lesions for which any non-malignant percutaneous tissue diagnosis is automatically considered discordant, resulting in the recommendation for repeat (usually surgical) biopsy.

Category 6: Known Biopsy - Proven Malignancy:

This category is reserved for examinations performed after biopsy proof of malignancy (imaging performed after percutaneous biopsy but prior to complete surgical excision) in which there are no mammographic abnormalities other than the known cancer that might need additional evaluation.

¹Mammography results are mandated to be reported using Final Assessment categories (Quality Mammography Standards: Final Rule. Federal Register. 1997;62:55988).

²Terminology in this table is reflective of the American College of Radiology (ACR). ACR-BI-RADS®--5th Edition. ACR Breast Imaging Reporting and Data System, Breast Imaging Atlas; BI-RADS. Reston VA. American College of Radiology, 2014. For more information, see www.acr.org. Reprinted with permission for the American College of Radiology. No other representation of this document is authorized without express, written permission from the American College of Radiology.

³The new BI-RADS® cut points for the risk of malignancy are as follows: 4A (>2% – ≤10%), 4B (>10% – ≤50%), 4C (>50% – <95%).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)

BSCR-C
4 OF 9



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

ULTRASOUND ASSESSMENT CATEGORY DEFINITIONS^{1,2}

BI-RADS® - ULTRASOUND FINDINGS

A. Assessment is Incomplete:

Category 0: Incomplete - Need Additional Imaging Evaluation:

There is a finding for which additional imaging evaluation is needed. This is almost always used in a screening situation. In this context, additional imaging evaluation includes the recording of (nonstandard) ultrasound images to supplement the standard images recorded for a screening examination. Note that this does not include repeat real-time scanning by the interpreting physician and/or colleague as long as additional images are not recorded. This respects the unique real-time nature of ultrasound and does not penalize its use.

Under certain circumstances, assessment category 0 may be used in a diagnostic ultrasound report, such as when equipment or personnel are not immediately available to perform a needed concurrent diagnostic mammography examination, or when the patient is unable or unwilling to wait for completion of a full diagnostic examination. Category 0 should not be used for diagnostic breast imaging findings that warrant further evaluation with MRI. Rather, the interpreting physician should issue a final assessment in a report that is made before the MRI examination is performed.

In most circumstances and when feasible, if a screening ultrasound examination is not assessed as negative or benign, the current examination should be compared to prior examination(s), if any exist. The interpreting physician should use judgment on how vigorously to attempt obtaining prior examinations, given the likelihood of success of such an endeavor and the likelihood that comparison will affect the final assessment. In this context, it is important to note that comparison to previous examination(s) may be irrelevant when a finding is inherently suspicious for malignancy.

Category 0 should be used for prior image comparison only when such comparison is required to make a final assessment. When category 0 is used in the context of awaiting prior examinations for comparison, there should be in place a tracking system guaranteeing with 100% reliability that a final assessment will be made within 30 days (preferably sooner), even if prior examinations do not become available. Some breast imaging practices may reasonably choose never to use category 0 in the context of awaiting prior examinations simply because they do not have a 100% reliable tracking system. If an ultrasound examination is assessed as category 0 in the context of awaiting prior examinations and then the prior examinations do become available, an addendum to the initial ultrasound report should be issued, including a revised assessment. For auditing purposes, the revised assessment should replace the initial assessment.

A need for previous studies to determine appropriate management might also temporarily defer a final assessment.

¹Mammography results are mandated to be reported using Final Assessment categories (Quality Mammography Standards: Final Rule. Federal Register. 1997;62:55988).

²Terminology in this table is reflective of the American College of Radiology (ACR). ACR-BI-RADS®--5th Edition. ACR Breast Imaging Reporting and Data System, Breast Imaging Atlas; BI-RADS. Reston VA. American College of Radiology, 2014. For more information, see www.acr.org. Reprinted with permission from the American College of Radiology. No other representation of this document is authorized without express, written permission from the American College of Radiology.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

ULTRASOUND ASSESSMENT CATEGORY DEFINITIONS^{1,2}

BI-RADS® - ULTRASOUND FINDINGS

B. Assessment is Complete — Final Categories:

Category 1: Negative:

There is nothing to comment on. This is a normal examination.

Category 2: Benign:

As with category 1, this is a “normal” assessment, but here the interpreter chooses to describe a benign finding in the ultrasound report. For example, the interpreter may choose to describe one or more simple cysts, intramammary lymph nodes, postsurgical fluid collections, breast implants, or complicated cysts/probable fibroadenomas that are unchanged for at least 2 or 3 years, while still concluding that there is no sonographic evidence of malignancy. On the other hand, the interpreter may choose not to describe such findings, in which case the examination should be assessed as negative (category 1).

Note that both category 1 and category 2 assessments indicate that there is no sonographic evidence of malignancy. Both should be followed by the management recommendation for routine age-appropriate screening. The difference is that category 2 should be used when describing one or more specific benign sonographic findings in the report, whereas category 1 should be used when no such findings are described (even if such findings are present).

¹Mammography results are mandated to be reported using Final Assessment categories (Quality Mammography Standards: Final Rule. Federal Register. 1997;62:55988).

²Terminology in this table is reflective of the American College of Radiology (ACR). ACR-BI-RADS®--5th Edition. ACR Breast Imaging Reporting and Data System, Breast Imaging Atlas; BI-RADS. Reston VA. American College of Radiology, 2014. For more information, see www.acr.org. Reprinted with permission from the American College of Radiology. No other representation of this document is authorized without express, written permission from the American College of Radiology.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

ULTRASOUND ASSESSMENT CATEGORY DEFINITIONS^{1,2}

BI-RADS® - ULTRASOUND FINDINGS

Category 3: Probably Benign:

Assessment category 3, probably benign, is not an indeterminate category for use simply when the radiologist is unsure whether to render a benign (BI-RADS® category 2) or suspicious (BI-RADS® category 4) assessment, but is one that is reserved for specific imaging findings known to have >0% but ≤2% likelihood of malignancy. For ultrasound, there is robust evidence that a solid mass with a circumscribed margin, oval shape, and parallel orientation (most commonly fibroadenoma) and an isolated complicated cyst have a likelihood of malignancy in the defined (≤2%), probably benign range, for which short-interval (6-month) follow-up sonography and then periodic sonographic surveillance may represent appropriate management. Similar data have been reported for clustered microcysts, but these data are less strong because they involve much fewer cases. The use of assessment category 3 for sonographic findings other than these three should be considered only if the radiologist has personal experience to justify a watchful-waiting approach, preferably involving observation of a sufficient number of cases of an additional sonographic finding to suggest a likelihood of malignancy within the defined (≤2%), probably benign range.

This edition of the BI-RADS® Atlas also emphasizes the recommendation that a category 3 assessment should not be made at screening; rather, this should be done only after completion of full diagnostic breast imaging examination. This recommendation is appropriate for screening mammography, for which batch interpretation usually is utilized, because in this setting there is no opportunity to complete the diagnostic workup before interpreting the screening examination. However, screening ultrasound almost always is interpreted online, so a full diagnostic examination also is completed while the patient remains in the breast imaging facility, and a single breast imaging report may be issued that combines the findings of both screening and diagnostic components of the examination. Hence, there is no purpose in recommending against category 3 assessment at screening ultrasound, because the diagnostic workup would be completed simultaneously. Note that for auditing purposes, the screening component of a category 3-assessed screening ultrasound examination will be audit-positive, not only because additional nonstandard (diagnostic) images will be recorded but also because a category 3 assessment at screening is defined as being audit-positive.

¹Mammography results are mandated to be reported using Final Assessment categories (Quality Mammography Standards: Final Rule. Federal Register. 1997;62:55988).

²Terminology in this table is reflective of the American College of Radiology (ACR). ACR-BI-RADS®--5th Edition. ACR Breast Imaging Reporting and Data System, Breast Imaging Atlas; BI-RADS. Reston VA. American College of Radiology, 2014. For more information, see www.acr.org.

Reprinted with permission from the American College of Radiology. No other representation of this document is authorized without express, written permission from the American College of Radiology.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

ULTRASOUND ASSESSMENT CATEGORY DEFINITIONS^{1,2}

BI-RADS® - ULTRASOUND FINDINGS

For category 3 assessments, the initial short-term follow-up interval is usually 6 months and involves the breast(s) containing the probably benign finding(s). Assuming stability at this 6-month examination, a category 3 assessment again is rendered with a management recommendation for a second short-interval follow-up examination in 6 months. Again assuming stability at this second short-interval follow-up, the examination is once more assessed as category 3, but now the recommended follow-up interval usually is lengthened to 1 year due to the already-observed 12-month stability. Note that although the 1-year follow-up coincides with the routine screening interval in the United States, a category 3 assessment is rendered to indicate that the period of imaging surveillance is still underway. As with surveillance using mammography, after 2 to 3 years of stability, the final assessment category should be changed to benign (BI-RADS® category 2). A benign evaluation may also be rendered before completion of category 3 analysis if, in the opinion of the interpreter, the finding has no chance of malignancy and is thus a category 2.

Category 4: Suspicious:

This category is reserved for findings that do not have the classic appearance of malignancy but are sufficiently suspicious to justify a recommendation for biopsy. The ceiling for category 3 assessment is a 2% likelihood of malignancy, and the floor for category 5 assessment is 95%, so category 4 assessments cover the wide range of likelihood of malignancy in between. Thus, almost all recommendations for breast interventional procedures will come from assessments made using this category. By subdividing category 4³ into 4A, 4B, and 4C, it is hoped that patients and referring clinicians will more readily make informed decisions on the ultimate course of action. An example of separating the BI-RADS® assessment category from the management recommendation occurs when a simple cyst, correctly assessed as BI-RADS® 2, undergoes cyst aspiration for pain control.

¹Mammography results are mandated to be reported using Final Assessment categories (Quality Mammography Standards: Final Rule. Federal Register. 1997;62:55988).

²Terminology in this table is reflective of the American College of Radiology (ACR). ACR-BI-RADS®--5th Edition. ACR Breast Imaging Reporting and Data System, Breast Imaging Atlas; BI-RADS. Reston VA. American College of Radiology, 2014. For more information, see www.acr.org. Reprinted with permission from the American College of Radiology. No other representation of this document is authorized without express, written permission from the American College of Radiology.

³The new BI-RADS® cut points for the risk of malignancy are as follows: 4A (>2% – ≤10%), 4B (>10% – ≤50%), 4C (>50% – <95%).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

ULTRASOUND ASSESSMENT CATEGORY DEFINITIONS^{1,2}

BI-RADS® - ULTRASOUND FINDINGS

Category 5: Highly Suggestive of Malignancy:

These assessments carry a very high probability (≥95%) of malignancy. This category initially was established to involve lesions for which 1-stage surgical treatment could be considered without preliminary biopsy in an era when preoperative wire localization was the primary breast interventional procedure. Nowadays, given the widespread acceptance of imaging-guided percutaneous biopsy, 1-stage surgery rarely, if ever, is performed. Rather, current oncologic management almost always involves tissue diagnosis of malignancy via percutaneous tissue sampling to facilitate treatment options, such as when sentinel node imaging is included in surgical management or when neoadjuvant chemotherapy is administered prior to surgery. Therefore, the current rationale for using a category 5 assessment is to identify lesions for which any nonmalignant percutaneous tissue diagnosis is considered discordant, resulting in the recommendation for repeat (usually vacuum-assisted or surgical) biopsy. Also note that whereas the fourth edition simply indicated that “appropriate action should be taken” as management for category 5 assessments, the fifth edition provides the more directed management recommendation that “biopsy should be performed in the absence of clinical contraindication.” This new text unequivocally specifies tissue diagnosis as the interpreting physician’s management recommendation for category 5 assessments, appropriately and effectively transferring the burden of establishing a contraindication to this recommendation to the referring clinician.

Category 6: Known Biopsy-Proven Malignancy:

This category is reserved for examinations performed after biopsy proof of malignancy (imaging performed after percutaneous biopsy but prior to surgical excision), in which there are no abnormalities other than the known cancer that might need additional evaluation.

¹Mammography results are mandated to be reported using Final Assessment categories (Quality Mammography Standards: Final Rule. Federal Register. 1997;62:55988).

²Terminology in this table is reflective of the American College of Radiology (ACR). ACR-BI-RADS®--5th Edition. ACR Breast Imaging Reporting and Data System, Breast Imaging Atlas; BI-RADS. Reston VA. American College of Radiology, 2014. For more information, see www.acr.org. Reprinted with permission from the American College of Radiology. No other representation of this document is authorized without express, written permission from the American College of Radiology.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

Discussion

This discussion is being updated to correspond with the newly updated algorithm. Last updated 05/03/17

NCCN Categories of Evidence and Consensus

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.

Table of Contents

Overview	1	Screening Recommendations for Women at Increased Risk	12
Breast Screening Components	3	Diagnostic Evaluation	15
<u>Clinical Encounter</u>	3	<u>Diagnostic Imaging After Screening Mammography Recall</u>	15
<u>Breast Cancer Risk Assessment</u>	4	<u>Diagnostic Mammography</u>	15
<u>Breast Imaging Modalities</u>	4	<u>Breast Ultrasonography</u>	16
Screening Recommendations for Women at Average Risk	7	<u>Diagnostic Breast MRI</u>	16
<u>Women with Average Risk Between the Ages of 25 and 39</u>	7	<u>Breast Tissue Biopsy</u>	17
<u>Women with Average Risk 40 Years and Older</u>	7	<u>Fine-Needle Aspiration (FNA) Biopsy</u>	17
		<u>Core Needle Biopsy</u>	17
		<u>Excisional Biopsy</u>	17
		<u>Diagnostic Evaluation For Symptomatic Findings on Physical Examination</u>	18
		<u>Palpable Mass in the Breast</u>	18
		<u>Women with Palpable Mass Aged 30 Years or Older</u>	18
		<u>Women with Palpable Mass Younger Than 30 Years of Age</u>	20
		<u>Nipple Discharge Without a Palpable Mass</u>	20
		<u>Asymmetric Thickening or Nodularity</u>	21
		<u>Skin Changes</u>	22
		<u>Breast Pain</u>	22
		Summary	23
		Table 1: Breast Cysts - Types and Definitions	24
		References	25



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

Overview

The average lifetime risk of breast cancer for a woman in the United States has been estimated at 12.3% (ie, 1 in 8 women).¹ For 2017, the American Cancer Society (ACS) estimates that 63,410 cases of female carcinoma in situ of the breast and 255,180 cases of invasive breast cancer (252,710 women and 2,470 men) will be diagnosed in the United States.² About 41,070 deaths are estimated for 2017.² The good news is that mortality rate from breast cancer has dropped 38% from 1989 through 2014.² This decrease has been partly attributed to mammographic screening.³

The National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology® (NCCN Guidelines®) for Breast Cancer Screening and Diagnosis are for facilitating clinical decision-making. The general public and health care providers need to be aware that mammography or any other imaging modality is not a stand-alone procedure. Neither the current technology of mammography or other imaging tests nor the subsequent interpretation of such tests is foolproof. Clinical judgment is needed to ensure appropriate management. The patient's concerns and physical findings must be taken into account along with imaging results and histologic assessment.

Breast Screening Components

Breast screening is performed in women without any signs or symptoms of breast cancer so that disease can be detected as early as possible, which allows early treatment to reduce the mortality and morbidity associated with the disease. A diagnostic breast evaluation differs from breast screening in that it is used to evaluate an existing problem (eg, palpable mass, discharge from the nipple).

The components of a breast screening evaluation are dependent on age and other factors such as medical and family history, and can include breast awareness (ie, patient familiarity with her breasts); regular clinical encounters, which include breast cancer risk assessment and clinical breast exam (CBE); breast imaging with screening mammography; and, in selected cases, ultrasound and breast MRI.

Clinical Encounter

The starting point of these guidelines for screening and evaluating breast abnormalities is a clinical encounter, which includes a complete medical history followed by breast cancer risk assessment and a CBE. The frequency of the clinical encounter depends on the age and risk assessment of the patient.

In a review of controlled trials and case-control studies that included CBE as part of the screening modality, sensitivity of CBE was found to be 54% and specificity 94%.⁴ Randomized trials comparing CBE versus no screening have not been performed. Rationale for recommending clinical encounter is to maximize earliest detection of breast cancers. Overdiagnosis and overtreatment is not a significant issue with CBE, as the majority of palpable cancers found on a CBE are invasive cancers. CBE is an important component of a clinical encounter and is important in order to detect early-stage palpable cancers, especially those that are mammographically occult (eg, lobular carcinomas). According to the NCCN panel, inspection of the breasts should be performed with the patient in both upright and supine positions. Positioning may be done so as to elicit any subtle shape or contour changes in the breast.⁴

Breast Awareness: Women should be familiar with their breasts and any changes to them.^{5,6} Data from a large randomized trial of breast self-examination (BSE) screening have shown that instruction in BSE



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

has no effect on reducing breast cancer mortality. In this study, 266,064 Chinese women not undergoing routine mammographic screening were randomized to either receive instruction in BSE or not.⁷ Compliance was encouraged through feedback and reinforcement sessions. After 10 to 11 years of follow-up, 135 breast cancer deaths in the instruction group and 131 in the control group were observed and the cumulative breast cancer mortality rates were not significantly different between the two arms (relative risk [RR], 1.04; 95% CI, 0.82–1.33; $P = .72$). The number of benign breast lesions detected in the BSE instruction group was higher than that detected in the control group. Nevertheless, women should be encouraged to be aware of their breasts since this may facilitate detection of interval cancers between routine screenings. The NCCN panel recommends breast awareness, specifically that all women should be familiar with their breasts and promptly report any changes to their health care provider.

Breast Cancer Risk Assessment

If the physical examination is negative in an asymptomatic woman, the next decision point is based on risk stratification. Women can be stratified into two basic categories for the purpose of screening recommendations: those at average risk and those at increased risk.

Risk assessment is outlined in the [NCCN Guidelines for Breast Cancer Risk Reduction](#). The increased risk category consists of six groups: 1) women with a prior history of breast cancer; 2) women ≥ 35 years of age with a 5-year risk of invasive breast cancer $\geq 1.7\%$ (per Gail Model); 3) women who have a lifetime risk $>20\%$ based on history of lobular carcinoma in situ (LCIS) or atypical ductal hyperplasia (ADH)/atypical lobular hyperplasia (ALH); 4) women who have a lifetime risk $>20\%$ as defined by models that are largely dependent on family history; 5) women between the ages 10 and 30 years with prior

thoracic RT (eg, mantle irradiation); and 6) women with a pedigree suggestive of or known genetic predisposition.

Breast Imaging Modalities

Screening Mammography

Of the various imaging modalities, mammography remains the most important as it is the only one to demonstrate a mortality reduction. A screening mammogram typically involves two x-ray images of each breast (ie, one taken from the top [craniocaudal] of the breast and the other from the side [mediolateral oblique]). Technical aspects of mammography can affect the quality of screening results. Digital mammography, which has replaced film-screen mammography in the United States, generates an electronic image of the breast and allows for computer storage and processing of the image, thereby increasing the ability to detect subtle abnormalities.^{8,9}

In a study of 49,528 women who underwent both film and digital mammography, no difference was seen in the overall accuracy of the two procedures.^{10,11} However, digital mammography was significantly more accurate in younger women with dense breasts, and there was a nonsignificant trend toward improved accuracy of film mammography in women aged 65 years and older. In another trial of women aged 45 to 69 years randomly assigned to film or digital screening mammography, the latter procedure was shown to result in a higher rate of cancer detection.¹²

More recently, combined use of digital mammography (two-dimensional, 2D) in conjunction with digital breast tomosynthesis (DBT) appears to improve cancer detection and reduce false-positive call-back rates.¹³⁻²² Tomosynthesis allows acquisition of three-dimensional (3D) data using a moving x-ray and digital detector. These data are reconstructed using computer algorithms to generate thin sections of images. The combined

use of 2D and DBT results in double the radiation exposure compared with mammography alone. However, this increase in radiation dose falls below dose limits of radiation set by the U.S. Food and Drug Administration (FDA) for standard mammography. The radiation dose can be minimized by newer tomosynthesis techniques that create a synthetic 2D image, which may obviate the need for a conventional digital image.^{14,23,24}

The presence of dense breast tissue decreases the sensitivity of mammography to detect small lesions and may obscure visualization of an underlying cancer. In addition, dense breast tissue as measured by mammography is increasingly recognized as an important risk factor for breast cancer.²⁵⁻²⁸ About half of all women of screening age have “dense” breast tissue referred to as “heterogeneously dense” or “extremely dense” by American College of Radiology (ACR) Breast Imaging Reporting and Data System (BI-RADS®) nomenclature. The presence of dense tissue is not abnormal and can change over time. Many states have passed legislation mandating patient notification of breast density, but few have required insurance coverage for supplemental screening.²⁹ However, the NCCN panel cautions that there is currently insufficient evidence to support routine universal supplemental screening in women with dense breasts and no other risk factors. Different supplemental imaging modalities may be considered based on risk and patient values/preference.³⁰

Screening Ultrasound

Due to limitations of mammographic screening, especially in women with dense breasts, other imaging modalities are being explored to supplement mammography, most commonly ultrasound and MR. Unlike mammographic screening, both technologies lack evidence from randomized controlled trials (RCTs) of screening efficacy, although ultrasound is widely used in the diagnostic setting. Most clinical

ultrasound screening studies have found increased cancer detection incremental to screening mammograms in women with dense breasts. For example, a large prospective study in women with dense breasts and elevated risk for breast cancer found that adding screening ultrasound to mammography identified an additional 4.3 cancers per 1000 women screened (95% CI, 1.1–7.2 cancers per 1000) but increased the number of false-positive results.³⁰ Subsequent follow-up studies showed similar results.^{31,32} However, in women with dense breasts, the mammographic sensitivity was found to be 50% (95% CI, 33.8%–66.2%) and the sensitivity of mammography plus ultrasound was 77.5% (95% CI, 61.6%–89.2%).³⁰ Application of screening ultrasound to women with dense breasts in clinical populations has produced similar results.³³

Although there is increasing evidence that breast ultrasonography can be useful in the incremental detection of breast cancer as an adjunct to screening mammography in the evaluation of women with dense breasts,^{30,31,34-36} the routine use of ultrasound as a universal supplemental *screening* test in women with average risk is *not* recommended by the NCCN panel at this time. Ultrasonography is commonly used for *diagnostic* follow-up of an abnormality seen on screening mammography and palpable clinical concerns.

Screening MRI

The sensitivity of contrast-enhanced breast MRI at detecting breast cancer is higher than the sensitivity of mammography, although the specificity of the former procedure is often lower, resulting in a higher rate of false-positive findings.³⁷ In addition, microcalcifications are not detectable with MRI.^{38,39} Similar to screening ultrasound, whether MRI screening impacts survival has not been addressed in randomized clinical trials. Therefore, careful patient selection for additional screening with MRI is needed. Although current evidence does not



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

support the use of breast MRI to screen women at average risk of breast cancer, the benefits of screening MRI for early detection of breast cancer in women with high risk, such as those ages 10 through 30 years with a history of prior thoracic radiation, a known genetic predisposition for breast cancer, or a strong family history of the disease have been demonstrated in multiple studies.⁴⁰⁻⁴⁸ The ACS has published guidelines recommending use of breast MRI as an adjunct to screening mammography in certain populations of women at high risk of breast cancer.⁴⁹ Nevertheless, a high false-positive rate for screening MRI was identified in several studies. For example, in one study of high-risk women, many of whom were young and had very dense breast tissue, screening MRI led to 3 times as many benign biopsies as mammography.⁵⁰

A single retrospective study of asymptomatic women with atypical hyperplasia or LCIS enrolled in a high-risk screening program has evaluated use of MRI in this population.⁵¹ Approximately half of the women underwent screening with mammography and MRI, whereas the other half was screened with mammography alone. For those undergoing both types of screening, MRI detected breast cancer in 4% of patients with LCIS who had negative mammogram results. MRI screening did not affect the rate of cancer detection in women with atypical hyperplasia. Women who underwent screening with MRI were more likely to be younger and premenopausal, and to have a stronger family history of breast cancer than those who were evaluated by mammography alone. However, only one woman with cancer detected by MRI following a negative mammography finding had reported a family history of breast cancer, and no difference was seen in the percentages of patients who ultimately developed cancer in the two groups.

The FDA has issued a safety alert stating that it is investigating the risks of brain deposits of gadolinium, the contrast agent used with breast

MRI. Studies have reported that deposits of gadolinium remain in the brain of some patients who undergo four or more contrast MRI scans, long after the last administration.⁵²⁻⁵⁴ In women with a history of thoracic radiation between ages 10 and 30 years, a known genetic predisposition to breast cancer, or a lifetime risk of >20% based on models such as Claus or Tyrer-Cuzick, based on current evidence, and considering the FDA warning (Gadolinium-based contrast agents), the NCCN panel continues to recommend an annual MRI as an adjunct to mammography. Women with LCIS/ALH/ADH should be considered for breast MRI based on emerging evidence of the benefits and their overall breast cancer risk.

Criteria for the performance/interpretation of high-quality breast MRI include a dedicated breast coil, radiologists experienced in breast MRI, and the ability to perform MRI-guided needle sampling and/or wire localization of MRI-detected findings. The ACR has published guidelines for the performance of contrast-enhanced MRI of the breast.⁵⁵

Other Breast Imaging Modalities

There is emerging evidence that breast scintigraphy and contrast-enhanced mammography may improve detection of early breast cancers among women with mammographically dense breasts;⁵⁶⁻⁵⁹ current evidence does *not* support their routine use as alternative screening procedures. Thermography and ductal lavage are *not* recommended by the NCCN panel for breast cancer screening or diagnosis. The FDA has issued a safety alert stating ductal lavage should not be a replacement for mammograms.⁶⁰



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

Screening Recommendations for Women at Average Risk

The NCCN panel recognizes that the primary purpose of screening women with average-risk for developing breast cancer is to detect breast cancer early, which allows treatment to decrease mortality and morbidity associated with breast cancer.

Women with Average Risk Between the Ages of 25 and 39:

The NCCN panel recommends a clinical encounter, which includes ongoing breast cancer risk assessment, risk reduction counseling, as well as a CBE every 1 to 3 years and encouraging women to be aware of their breasts and promptly report any changes to their health care provider.

Although the screening CBE by itself does not rule out disease, the high specificity of certain abnormal findings by highly qualified clinicians increases the probability of finding certain breast cancers (eg, lobular carcinoma). The NCCN panel believes that a clinical encounter provides an opportunity for providers to perform a CBE, conduct a breast cancer risk assessment, provide risk reduction recommendations, and counsel on healthy lifestyles.

Women with Average Risk 40 Years and Older:

The NCCN panel recommends annual clinical encounter, which includes ongoing breast cancer risk assessment, risk reduction counseling, as well as a CBE, and encourages women to be aware of their breasts and promptly report any changes and annual screening mammography (category 1 recommendation) with the *consideration* of tomosynthesis. Women electing to undergo screening mammography should be counseled regarding its potential benefits, risks, and limitations. The NCCN panel is in agreement with ACS and other organizations that annual screening mammograms in average-risk

women age 40 years and older should be covered by the health care payers without additional cost-sharing or copayments.

Mammographic screening and subsequent treatment has been shown to decrease breast cancer mortality beginning at age 40.^{61,62} Meta-analysis of invitational RCTs, observational studies, and computer modeling of mammographic screening consistently show benefit, although the magnitude of benefit has varied in part due to the diversity of study designs and screening frequency. However, the RCTs are now old and may not reflect current mammography technology, interpretation, and oncologic care. Therefore, effectiveness may be better estimated in more modern observational studies.

The mammography screening guidelines put forth by various organizations vary with respect to age to initiate screening, the frequency of screening, and when to stop screening.^{61,62} The assessment of the benefits of mammography versus the risks based on age are weighed on different scales by different organizations.

The NCCN panel continues to support its long-standing recommendation of *annual* screening mammography beginning at age 40 (category 1 recommendation), as it results in the greatest mortality reduction, most lives saved, and most life years gained.

The NCCN panel has not established an upper age limit for screening. According to the panel, if a patient has severe comorbid conditions limiting her life expectancy and no further intervention would occur based on the screening findings, then the patient should not undergo screening, regardless of her age.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

Rationale for Mammographic Screening Starting at Age 40:

Reduction in breast cancer-related mortality is the major benefit of mammographic screening for breast cancer. This benefit is evident across studies, including RCTs, case-controlled observational studies, and computer modelling studies.

While breast cancer screening guidelines put forth by all the organizations acknowledge mortality reduction benefit from current studies of mammography screening in women 40 to 49 years of age, those recommending breast cancer screening to begin at age 50⁶² view the benefits of screening as being balanced by the harms of screening during this decade. Other organizations, who have recommended screening commencement at age 45 as a “strong” recommendation, have shown the absolute benefit of ages 45 to 49 to be very similar to ages 50 to 54.⁶¹ While showing there is benefit of screening for ages 40 to 44, a “qualified” rather than a “strong” recommendation is given for the younger age group due to the lower absolute benefit. However, the “qualified” recommendation means “most” women would want the earlier screening and only a “small proportion” would not.⁶¹

Benefits of Mammographic Screening:

Systematic reviews of RCTs have generally shown a reduction in breast cancer mortality with mammography screening.⁶³

The UK Age trial specifically studied the effect of film-screen mammographic screening starting at age 40 years.⁶⁴ A mean of 10.7 years of follow-up showed a non-statistically significant breast cancer mortality reduction in women invited to screening (RR, 0.83; 95% CI, 0.66–1.04).⁶⁴ A follow-up of the UK AGE trial was carried out to study breast cancer mortality and incidence at a median of 17.7 years of follow-up, an increase of 7 years from the previous analysis.⁶⁵ There continued to be a non-significant overall reduction in risk of breast

cancer mortality (RR, 0.88; 95% CI, 0.74–1.04) during a median of 17 years of follow-up. However, the reduction in breast cancer mortality noted in the first 10 years after diagnosis was now significant in the group that underwent screening compared with the control group (RR, 0.75, 0.58–0.97).⁶⁵ Other trials included women who were up to age 49 years at the time of entry into the trial, who were therefore in their 50s during the screening intervention. The results of the UK Age trial support the importance of annual mammography screening in women ages 40 to 49 years of age to reduce breast cancer-related mortality.⁶⁵

A Swedish study compared breast cancer mortality rates in women 40 to 49 years living in different counties. Counties included those which invited women for screening starting at age 40 and others that did not invite the women to be screened at age 40 and started screening at age 50.⁶⁶ After an average 16 years of follow-up, the investigators observed an overall 29% mortality reduction (RR, 0.71; 95% CI, 0.62–0.80). For age groups 40 to 44 and 45 to 59 years, the RR estimates were 0.82 (95% CI, 0.67–1.00) and 0.63 (95% CI, 0.54–0.75).⁶⁶ Although the estimated reduction in breast cancer mortality was smaller for ages 40 to 44 compared with ages 45 to 49, the reduction in mortality seen for ages 40 to 44 was **observational studies**

still substantial.⁶⁶

It is important to note that the RCTs studying the benefits of screening mammography used screen film mammography, sometimes using only a single view. Therefore, they may not reflect results obtained with modern advances in imaging. Digital mammography has been shown to detect more breast cancers in women with dense breasts, which is common in younger women. The more recent observational studies better quantify the effectiveness of screening in the context of improved imaging techniques.

NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

Case-control observational studies have shown benefits of reduction in breast cancer mortality ranging from 40% to 45%.^{67,68} A meta-analysis of observational case-control studies found a significant reduction in breast cancer mortality with mammographic screening for women aged 40 to over 79 years of age with a 48% mortality reduction (odds ratio [OR] 0.52; 95% CI, 0.42–0.65) after adjustment for self-selection.⁶⁹

Relevant to the North American population, data from a Canadian study showed a mortality reduction of 44% (CI, 33%–55%) among screened women ages 40 to 49 years, which was similar to the overall reduction in mortality of 40% (CI, 33%–48%) found among women ages 40 to 79 years.⁶⁸

A retrospective analysis evaluating the benefits of mammographic screening of women aged 40 to 49 years found that mammography-detected breast cancer coincides with lower-stage disease at detection, resulting in reduced treatment morbidity and lower rates of recurrence.⁷⁰ A population-based study of data from the Netherlands Cancer Registry estimated the impact of tumor size in women with breast cancer in two time intervals: 1999 to 2005 and 2006 to 2012. The year 2005 was used to divide the data two time intervals studies, because trastuzumab and other effective adjuvant therapy were introduced after this year in the Netherlands. The analysis found tumor size remained a critical component of survival even with the availability of new and effective systemic therapy options.⁷¹ These findings reiterate that fact that diagnosing breast cancer at an early stage is important.

The Cancer Intervention and Surveillance Modeling Network (CISNET) models from 2009 demonstrate a 29% to 54% (mean 39%) mortality reduction for annual screening for women ages 40 to 84 years.⁷² The CISNET models from 2015 based on digital screening mammography show greater mortality reduction benefit.⁷³ Benefits for screening younger women (in their 40s) are more favorable when considered from

the perspective of years of life saved compared exclusively to mortality reduction.⁷⁴ Women in their 40s have the highest number of life years at risk to be lost due to longevity even though their breast cancer risk is smaller. Breast cancer is the leading cause of cancer deaths for women in their 40s.

Women should be informed of the evidence demonstrating the value of detecting breast cancer early, before symptoms develop. The benefits of early detection include mortality reduction, less aggressive treatment, and a wide range of treatment options.

Harms of Mammographic Screening:

The harms or risk profile for breast cancer screening is weighted differently by different organizations.^{61,62} This is a very subjective rating as there are limited data regarding a woman's perspective of the harms of screening. The clinical practice guidelines that recommend delaying screening to age 50 and older⁶¹ place a greater emphasis on the risks of screening mammography, specifically false-positive results and overdiagnosis. The reduction in breast cancer mortality is valued highly by most women, whereas many women do not consider false positives and potential overdiagnosis to be a "harm."⁷⁵ In this study, 63% of women thought 500 or more false positives per life saved was acceptable.⁷⁵

The NCCN panel believes that the harms analysis of mammographic screening is most informative if it includes the net harms of mammographic screening in individuals who underwent screening versus those who did not. According to the NCCN panel, the major harm related to *not performing* any screening for breast cancer is diagnosis of later-stage breast cancer, which may require more extensive therapy and may prove lethal. There is evidence showing that women diagnosed with breast cancer who did not undergo screening



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

had substantially more need for chemotherapy and more extensive surgery than women who underwent routine screening.⁷⁶

Furthermore, absence of mammographic screening for breast cancer does not mean absence of breast-related problems. Non-screened women develop signs and symptoms leading to diagnostic investigation, false-positive biopsies, or potential diagnosis of non-lethal conditions.

A mammogram result is often considered a false positive when it prompts additional imaging tests and/or biopsy in an abnormality that is not cancerous. False-positive results can occur at any age. It is important to distinguish between recalls from screening and false-positive recommendations for biopsy. Recalls are defined by the FDA as “incomplete” and not positive. Recalls are resolved by obtaining incremental diagnostic mammographic imaging or ultrasound with the vast majority of recalls proving negative and not requiring biopsy. The frequency of recalls from screening are the same per decade whether screening begins at age 40 or age 50.⁶² While recalls are commonly thought to be higher in younger women, this primarily reflects higher recall rates at the prevalent or initial screen when prior mammograms are not available for comparison and not at the age at which screening commences. The initiation of screening mammography at age 50 would shift this “prevalent” false positive to that decade. Furthermore, the decade-long false-positive biopsy recommendation rate is actually somewhat lower when screening begins at age 40 compared to age 50. Less than 1% of screened women per year will be recommended for a biopsy that proves benign, whether annual screening commences at age 40 or 50. The vast majority of false-positive biopsies are now performed as outpatient image-guided needle biopsies using local anesthesia.

Those considering false positives as one of the harms of screening note psychosocial consequence as one of the negative consequences of false positives.⁷⁷ However, a cross-sectional survey of women’s attitudes toward false positives found that women consider false positives as an acceptable consequence.⁷⁵

Overdiagnosis is the detection of a condition by screening that would not have become apparent by usual care absent screening. Overdiagnosis may lead to overtreatment, which is the more significant problem. It is important to understand that overdiagnosis would not influence the age to initiate screening or the screening interval. The mammographic abnormality that leads to a potential overdiagnosis does not go away without treatment. If the age to initiate screening is raised from 40 to 45 years or 50 years or the screening interval were lengthened to biennial, the potential overdiagnosis would occur at the next mammogram that showed the imaging abnormality.

Overdiagnosis is difficult to measure because neither the clinician, pathologist, nor the patient can be sure whether the abnormality detected by screening would be harmless or life threatening to the patient. Furthermore, overdiagnosis assumes the level or amount of diagnosis by symptomatic usual care is optimal. The estimates of overdiagnosis vary widely between various studies (from almost none up to 54%^{61,63,78-80}) due to methods and parameters used for estimation and whether ductal carcinoma in situ (DCIS) is included or excluded. Furthermore, overdiagnosis estimates vary by age and duration of follow-up.

The most reliable estimates of overdiagnosis would be from RCTs in which there was no formal screening offered to the control group for a long period at the end of the screening period. The Malmö randomized trial, in which the older-age invited cohort group was not routinely



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

screened at the end of the trial,⁸¹ showed after an average of 15 years follow-up an overdiagnosis rate of 10%, which included invasive cancer and DCIS. The rate was 7% for invasive cancer.⁸¹ The National Breast Screening Studies in Canada conducted two randomized trials that included a control group that did not receive routine screening at the end of the trial. The follow-up period was 13 years. In the first trial, in which women were aged 40 to 49 years at recruitment, the estimated overdiagnosis was 14%. In the second trial, in which women were aged 50 to 59 years at recruitment, the estimated overdiagnosis rate was 11%.^{82,83} Using these 3 studies, the UK review estimated overdiagnosis (including DCIS) to be 10.7%.⁸⁴ However, analysis of the UK AGE trial, which included women aged 40 to 49 years, showed a very low rate of overdiagnosis of 1%,⁸⁵ a value similar to estimates from Sweden for women in their 40s.⁶⁶ A recently reported population-based screening study showed a rate of only 0.3% overdiagnosis after 12 years of follow-up in either invited or uninvited women (n = 988, 090) and a 46% reduction in breast cancer mortality among attenders.⁸⁶

Prevention of cancer death is highly valued compared with false-positive results/overdiagnosis by most women.⁷⁵ Science cannot predict which breast cancer may be overdiagnosed or be potentially lethal in any one individual. Personalized treatment programs are recommended. The treatment of cancer may cause suffering and anxiety, but that suffering is likely worth the gain from the potential reduction in breast cancer mortality. According to the NCCN panel, the risk of overdiagnosis and false positives are outweighed by the benefit of mortality reduction in determining the age to recommend starting screening.

The NCCN panel emphasizes adopting strategies and research to reduce the harms of screening (false positives and overdiagnosis) rather than raising the age to initiate screening to potentially delay these

issues. This includes newer imaging modalities that improve the detection of breast cancer with fewer recalls (eg, tomosynthesis). Research to better define the biology of breast cancer is needed so that lesions that are not destined to progress are either not treated or are treated less aggressively.

Screening Interval and Rationale for Annual Mammogram Screening:

Another consideration is the time interval between screening exams. Performing screening mammography annually versus every other year remains controversial. Most studies and models suggest incremental benefit with annual screening, especially among younger women and premenopausal women.^{61,62,72,87} The evaluation of benefits versus risk strongly supports the value of screening and the importance of adhering to a schedule of regular mammograms.

The NCCN panel believes that the benefits of annual mammography outweigh the risks. Breast cancer mortality is estimated to be lower with annual compared to biennial screening mammograms.⁷² Additionally, mammograms can often detect a lesion 2 years before the lesion is discovered by CBE. Interval cancer rates are lower among annually screened women. To reduce mortality from breast cancer, yearly screening is thought to be more beneficial. The panel also acknowledges that incomplete compliance will alter the outcome of any recommendation.

An evaluation of the CISNET modeling of benefits of screening women between 40 to 49 years found that using *annual* digital mammography saves 30% more lives and 34% more life-years than *biennial* digital mammography.⁸⁸ Also, with annual digital screening mammography, the deaths averted (0.6/1000) are similar for ages 40 to 44 and 45 to 49 (0.7/1000).^{87,89}

NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

A decline in breast cancer specific-mortality was observed in a cohort of women for every additional annual mammogram performed 5 years prior to breast cancer diagnosis; this further emphasizes the importance of annual mammography.⁹⁰ The results of a primary analysis to estimate the association between incidence of DCIS detected by screening and subsequent invasive interval cancer incidence showed a DCIS detection rate of 1.5 per 1000 screened and a reduction of one invasive interval cancer per 1.5 to 3 DCIS cases detected.⁹¹

While the risk of false positives are greater with annual compared to biennial mammograms,⁶² the panel believes that the lower mortality and morbidity of annual screening outweighs this harm.

Age to Stop Mammographic Screening:

There are limited RCT data regarding screening of elderly women, because most trials for breast screening have used a cutoff age of 65 or 70 years.⁹²⁻⁹⁴ However, observational studies and computer models show mortality benefit to age 80 to 84.^{61,72} Considering the high incidence of breast cancer in the elderly population, the screening guidelines used for women who are age 40 or older are recommended in the elderly as well. Clinicians should always use judgment when applying screening guidelines. The mortality benefit of screening mammography is often delayed for 5 to 7 years in RCTs that emphasize the importance of life expectancy and overall health when considering age to stop screening. Mammography screening should be individualized weighing its potential benefits/risks in the context of the patient's overall health and estimated longevity.⁹⁵ If a patient has severe comorbid conditions limiting her life expectancy and no intervention would occur based on the screening findings, then the patient should not undergo screening, regardless of her age.^{95,96}

Screening Recommendations for Women at Increased Risk

Women with Prior History of Breast Cancer: These women are treated according to the recommendations outlined in [NCCN Guidelines for Breast Cancer](#).

Women Aged 35 Years or Older with a 5-Year Risk of Invasive Breast Carcinoma Greater Than or Equal to 1.7% by the Modified Gail Model: For women aged 35 years and older, a risk assessment tool is available to identify those who are at increased risk. The National Cancer Institute (NCI) and the National Surgical Adjuvant Breast and Bowel Project (NSABP) Biostatistics Center has developed a computerized interactive risk-assessment tool based on the modified Gail model⁹⁷⁻¹⁰¹ that can be accessed at: <http://www.cancer.gov/bcrisktool/Default.aspx>, which provides risk projections on the basis of several risk factors for breast cancer. The modified Gail model assesses the risk of invasive breast cancer as a function of age, menarche, age at first live birth or nulliparity, number of first-degree relatives with breast cancer, number of previous benign breast biopsies, atypical hyperplasia in a previous breast biopsy, and race. The model calculates 5-year and lifetime projected probabilities of developing invasive breast cancer and can be used to identify women who are at increased risk. The Gail model should not be used for women with a predisposing gene mutation, a strong family history of breast or ovarian cancer suggestive of a genetic predisposition, women with a prior history of thoracic radiation, or for those with LCIS.

The Gail model was updated using combined data from the Women's Contraceptive and Reproductive Experiences (CARE) study and the SEER database, as well as causes of death from the National Center for Health Statistics, to provide a more accurate determination of risk for African-American women.¹⁰² It has also been updated using the data



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

from the Asian American Breast Cancer Study (AABCS) and the SEER database to provide a more accurate risk assessment for Asian and Pacific Islander women in the United States.¹⁰³

Increased risk of developing breast cancer is defined by the modified Gail model for women ≥ 35 years of age as a 5-year risk of 1.7% or greater. This is the average risk of a 60-year-old woman, which is the median age of diagnosis of breast cancer in the United States. The 5-year predicted risk of breast cancer required to enter the NSABP Breast Cancer Prevention Trial of tamoxifen versus placebo, as well as the Study of Tamoxifen and Raloxifene (STAR) trial, was 1.7% or greater. As previously mentioned, the modified Gail model risk assessment tool also provides an estimate of a woman's lifetime risk of breast cancer. However, this estimate is based on the Gail model risk criteria, which differ from criteria used in risk assessment models predominantly based on family history (see below). Lifetime breast cancer risk as determined by the Gail model is not used in these guidelines to determine whether a woman is eligible for screening breast MRI.

For a woman aged 35 years or older with a 5-year risk $\geq 1.7\%$, the NCCN panel encourages breast awareness and recommends a clinical encounter every 6 to 12 months and annual digital mammography, with the consideration of tomosynthesis, to begin at the age identified as being at increased risk by the Gail model. In addition, according to the NCCN panel, women in this group should be counseled for consideration of risk-reduction strategies in accordance with the [NCCN Guidelines for Breast Cancer Risk Reduction](#).

Women Who Have a Lifetime Risk $>20\%$ Based on History of LCIS or ADH/ALH: A diagnosis of LCIS or ADH/ALH is associated with high risk of development of cancer in either breast.¹⁰⁴⁻¹⁰⁶

For women with a history of LCIS or ADH/ALH, the NCCN panel encourages breast awareness and recommends a clinical encounter every 6 to 12 months beginning at the age of diagnosis and annual digital mammography, with the consideration of tomosynthesis, beginning at the age of diagnosis of LCIS or ADH/ALH but not less than 30 years of age. In addition, according to the NCCN panel, annual MRI should be considered beginning at the age of diagnosis of LCIS or ADH/ALH but not less than age 25 (based on emerging evidence).⁵¹ Women in these groups should also be considered for risk reduction strategies in accordance with the [NCCN Guidelines for Breast Cancer Risk Reduction](#).

Women with a Lifetime Risk of Breast Cancer $>20\%$ Based on Models Largely Dependent on Family History: A lifetime risk of breast cancer of $>20\%$ as assessed by models based largely on family history is another risk threshold used in the guidelines to identify a woman as a potential candidate for risk reduction strategies, as well as to direct screening strategies. According to the ACS guidelines for breast screening, MRI may be performed as an adjunct to mammography⁴⁹ in a high-risk woman if her lifetime risk of breast cancer is approximately 20% or greater based on models that rely mainly on family history. A cancer genetic professional should be involved in determining the lifetime risk of the individual based on models dependent on family history. These include Claus,¹⁰⁷ Tyrer-Cuzick,¹⁰⁸ and other models.¹⁰⁹⁻¹¹¹ BRCAPRO¹¹² and Breast and Ovarian Analysis of Disease Incidence and Carrier Estimation Algorithm (BOADICEA)¹¹³ are more commonly used to estimate the risk of *BRCA* mutations. Strong genetic association between breast and ovarian cancer has been demonstrated in some families by linkage analyses.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

For a woman with a >20% lifetime risk of breast cancer based on models largely dependent on family history, the NCCN panel encourages breast awareness and clinical encounter every 6 to 12 months to begin at the age identified as being at increased risk. The NCCN panel recommends annual digital mammography, with the consideration of tomosynthesis starting from 10 years prior to the youngest family member but not less than age 30. In addition, in accordance with the ACS guidelines,⁴⁹ the NCCN panel recommends annual breast MRI to begin 10 years prior to the youngest family member diagnosed but not less than 25 years of age for women who have a lifetime risk of breast cancer >20% based on models that rely mainly on family history. According to the NCCN panel, women in this group should be asked to consider risk reduction strategies in accordance with the [NCCN Guidelines for Breast Cancer Risk Reduction](#).

Women Who Have Received Prior Thoracic Irradiation Between the Ages of 10 to 30 Years: Results from several studies have demonstrated that women who received thoracic irradiation in their second or third decade of life have a substantially increased risk of developing breast cancer by age 40 years.¹¹⁴⁻¹¹⁹ For example, in the Late Effects Study Group trial, the overall risk of breast cancer associated with prior thoracic irradiation at a young age was found to be 56.7-fold (55.5-fold for female patients) greater than the risk of breast cancer in the general population.^{115,118} The RR of female breast cancer according to follow-up interval was 0 at 5 to 9 years; 71.3 at 10 to 14 years; 90.8 at 15 to 19 years; 50.9 at 20 to 24 years; 41.2 at 25 to 29 years; and 24.5 at >29 years.¹¹⁸ Results from a case-control study of women treated with thoracic radiation at a young age for Hodgkin lymphoma indicated that the estimated cumulative absolute risk of breast cancer at 55 years of age was 29.0% (95% CI, 20.2%–40.1%)

for a woman treated at 25 years of age with at least 40 Gy of radiation and no alkylating agents.¹²⁰ Although there is a concern that the cumulative radiation exposure from mammography in a young woman may itself pose a risk for cancer, it is felt that the additional radiation in this population is negligible compared to overall radiation exposure. Findings from a survey of breast screening practices in this population of patients suggest that a sizable segment of this group is not undergoing regular mammographic screening.¹²¹

For women aged 25 years and older who have received prior thoracic irradiation, the NCCN panel recommends encouraging breast awareness, and a clinical encounter every 6 to 12 months be initiated 8 to 10 years after radiation exposure.¹²² Breast imaging assessments with annual digital mammograms, with the consideration of tomosynthesis, and annual MRI as an adjunct to mammograms⁴⁰ are recommended to begin 8 to 10 years after radiation exposure in those aged 25 years or older.

For women younger than 25 years who have received prior thoracic irradiation, the NCCN panel recommends encouraging breast awareness, counseling on risk, and an annual clinical encounter starting 8 to 10 years after radiation therapy.

Women with a Pedigree Suggestive Of or With a Known Genetic Predisposition: Accurate family history information is needed to adequately assess a woman's breast cancer risk. Familial cancers share some but not all features of hereditary cancers. For example, although familial breast cancers occur in a given family more frequently than expected based on statistics, they generally do not exhibit inheritance patterns or onset age consistent with hereditary cancers. Familial breast cancers may be associated with chance clustering,



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

genetic variations in lower-penetrance genes, a shared environment, small family size, and/or other factors.

The NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast and Ovarian include recommendations for referral to a cancer genetics professional for further evaluation for individuals who have either a personal history or a close family history meeting certain criteria and also list screening recommendations for common hereditary syndromes that confer increased risk for breast and ovarian cancer. (See [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast and Ovarian](#)).

Diagnostic Evaluation

Breast symptoms are common among women. A retrospective study of women aged 40 to 70 years showed that 16% (total visits of 23 per 100 women) of women will present with symptoms to their provider during a decade with higher frequency among women ages 40 to 59 years compared to older women.¹²³ Pain is found to be the most common symptom followed by palpable mass. In addition, palpable areas of concern are identified during a breast physical exam. Breast clinical findings are not specific and there is variability in interpretation. Each symptom is associated with a risk of malignancy and warrants diagnostic evaluation even though most symptoms will be determined to be benign in etiology. Women younger than age 40, who are not usually recommended for routine breast screening, also frequently present for breast symptoms.

Unlike imaging for screening, which is used to detect cancer in asymptomatic women, diagnostic evaluation is used to characterize a clinical finding or possible abnormality found during screening. There is confusion regarding the term “diagnostic” imaging, as it is applied to two very different situations: 1) imaging for clinical finding such as a

palpable mass; and 2) incremental imaging after a possible abnormal screening mammogram in an asymptomatic woman (also referred to as recall or call-back). To add further confusion, insurance carriers may consider a routine mammogram to be “diagnostic” in certain asymptomatic women (eg, in women with prior cancer). Diagnostic evaluation in this review will be restricted to the former two situations.

Diagnostic evaluation may include physical examination and diagnostic imaging for symptomatic women and diagnostic imaging for women recalled from screening. Diagnostic imaging may include diagnostic mammography, ultrasonography, and at times diagnostic breast MRI. The eventual decision regarding need for tissue sampling is based on level of suspicion on imaging and/or clinical examination. Biopsy is needed in situations where imaging is negative but clinical findings are suspicious since imaging is not completely sensitive for cancer detection.

While the term “diagnostic” implies diagnosis, imaging results are often not specific enough to be truly “diagnostic.”

Diagnostic Imaging After Screening Mammography Recall

Diagnostic Mammography

Screening mammography consists of two standard x-ray images of each breast, whereas a diagnostic mammogram includes additional views, such as spot compression views or magnifications views, to investigate the finding in question. Diagnostic mammography is associated with higher sensitivity but lower specificity as compared to screening mammography. DBT may replace traditional diagnostic mammographic imaging in certain situations.¹²⁴⁻¹²⁶

Frequently, especially for masses or asymmetries, diagnostic ultrasound is also performed. Each imaging modality may be positive



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

or negative, which allows four outcomes: both imaging modality results are negative; both are positive; mammogram is positive and ultrasound is negative; and mammogram is negative and ultrasound is positive. In general, a “final” combined imaging assessment category is rendered after a “recall” from screening, which is the most suspicious imaging outcome assessment.

The mammographic final assessments are mandated by MQSA and are reported using the similar ACR BI-RADS® assessment categories, which classify likelihood of the breast findings into six final assessment categories.¹²⁷ The BI-RADS® assessment categories help to standardize both the reporting of mammographic findings and the recommendations for further management. The definitions of the mammogram assessment categories are outlined in *Mammographic Assessment Category Definitions* in the algorithm.

NCCN Recommendations for Mammogram BI-RADS® Assessment Categories 1, 2, 3, 4, 5, and 6

For BI-RADS® category 1 (negative finding) or category 2 (benign), the panel recommends resuming routine screening.

For BI-RADS® category 3 (probably benign), the panel recommends diagnostic mammograms at 6 months, then every 6 to 12 months for 1 to 2 years as appropriate. If the lesion remains stable or resolves mammographically, the patient resumes routine screening intervals for mammography. If, in any of the interval mammograms, the lesion increases in size or changes its benign characteristics, a biopsy is then performed. The exception to this approach of short-term follow-up is when a return visit is uncertain or the patient strongly desires or has a strong family history of breast cancer. In those cases, initial biopsy with histologic sampling may be a reasonable option.

For BI-RADS® Categories 4 and 5, tissue diagnosis using core needle biopsy (preferred) or needle localization excisional biopsy with specimen radiograph is necessary. When a needle biopsy (aspiration or core needle biopsy) is performed, concordance between the pathology report and the imaging finding must be obtained.^{128,129} For example, a negative needle biopsy associated with a spiculated category 5 mass is discordant and clearly would not be an acceptable diagnosis. When the pathology and the imaging are discordant, the breast imaging should be repeated and/or additional tissue sampled or excised; surgical excision is recommended when pathology/image remain discordant. Women with a benign result exhibiting pathology/image concordance should be followed with mammography every 6 to 12 months for 1 to 2 years before returning to routine screening.

For BI-RADS® category 6 (proven malignancy), the patient should be managed according to the [NCCN Guidelines for Breast Cancer](#).

Breast Ultrasonography

Imaging by ultrasound is an important adjunct for diagnosing breast cancer.¹³⁰ However, breast ultrasonography does not detect most microcalcifications.^{30,41,131-133} The definitions of the ultrasound assessment categories are outlined in *Ultrasonographic Assessment Category Definitions* in the algorithm.

Diagnostic Breast MRI

MRI can also play a role in the diagnostic setting. For patients with skin changes consistent with serious breast disease, consideration of breast MRI is included in the guidelines for those with benign biopsy of skin or nipple following BI-RADS® category 1-3 assessment. Since a benign skin punch biopsy in a patient with a clinical suspicion of inflammatory breast cancer (IBC) does not rule out malignancy, further evaluation is recommended. There is evidence that certain MRI features may



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

facilitate diagnosis of IBC.¹³⁴ MRI may be used for suspicious nipple discharge when mammography and ultrasound are not diagnostic.¹³⁵⁻¹³⁷

Breast Tissue Biopsy

Breast biopsy is recommended if diagnostic imaging findings or clinical findings are suspicious or highly suggestive of malignancy.

Fine-Needle Aspiration (FNA) Biopsy

An FNA biopsy involves use of a smaller-bore needle to obtain cytologic samples from a breast mass. Advantages of FNA biopsy include its minimally invasive methodology and low cost,^{138,139} whereas the need for pathologists with specific expertise in the interpretation of test results and the necessity of performing a follow-up tissue biopsy when atypia or malignancy is identified are disadvantages of the procedure. FNA of nonpalpable lesions can be performed under imaging guidance (eg, ultrasound), although there is evidence to indicate that both core needle biopsy and excisional biopsy are more accurate than FNA in the evaluation of nonpalpable breast lesions.^{140,141}

Core Needle Biopsy

A core needle biopsy, also called percutaneous core breast biopsy, is a procedure that typically involves obtaining multiple cores of solid tissue using standard techniques.^{142,143} It can be performed under imaging guidance (eg, stereotactic [mammographic] ultrasound or MRI) or directed by palpation. Advantages of breast core needle biopsy include increased accuracy over FNA when the procedure is performed in situations where no mass is palpable and an ability to obtain tissue samples of sufficient size so as to eliminate the need for a follow-up biopsy to confirm malignancy.¹⁴⁴ In some situations, the core needle biopsy is performed under vacuum assistance, which can facilitate collection of adequate tissue from a breast lesion without the need for multiple needle insertions.¹⁴⁵⁻¹⁴⁷ Marker clip placement is done at the

time of core needle biopsy so that the radiologist can identify the location of the lesion in the event that it is entirely removed or disappears during neoadjuvant treatment of a breast cancer.¹⁴⁸ With a few exceptions, core needle biopsy is preferred in the NCCN Guidelines over surgical excision when tissue biopsy is required. Sensitivity for core needle biopsy directed by ultrasound or stereotaxis is 97% to 99%.⁸⁹ According to the NCCN panel, surgical excision is appropriate if unable to perform core needle biopsy.

Excisional Biopsy

An excisional biopsy involves removal of the entire breast mass or suspicious area of the breast by a surgeon in an operating room setting. Needle or wire localization is done by the radiologist immediately prior to an excisional biopsy of a nonpalpable mammographic or sonographic finding to direct surgical excision. The wire localization may bracket a lesion that had a clip placed in it at the time of the core needle biopsy.¹⁴⁸ Newer localization methods using radionuclide seeds or reflector devices are being explored.

Excisional biopsy is included in the NCCN Guidelines as an option when tissue biopsy is required. Although excisional biopsy is a more invasive method than core needle biopsy and requires needle localization when lesions are not palpable, there are situations where larger tissue samples may be needed. In most cases, excisional biopsy is recommended following diagnosis by core biopsy of an indeterminate lesion, atypical hyperplasia, LCIS, or a benign and image-discordant lesion. Other histologies that may require additional tissue include mucin-producing lesions, potential phyllodes tumor, papillary lesions, radial scars, or other histologies of concern to the pathologist.^{139,144,149,150} Support for this recommendation includes results of studies demonstrating an underestimation of cancer when atypical hyperplasia and LCIS are diagnosed by core needle biopsy.¹⁵¹⁻¹⁵⁶ However, there



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

are situations (eg, select cases of LCIS, ALH, papillomas, fibroepithelial lesions, and radial scars) where close observation may be substituted for excisional biopsy in select patients.^{139,149,157-163}

Diagnostic Evaluation For Symptomatic Findings on Physical Examination

In general, the breast imaging evaluations after physical exam include mammography and ultrasound. The addition of ultrasound to diagnostic mammography significantly increases cancer detection and detection of specific benign findings such as cysts. Imaging for women younger than age 30 begins with ultrasound while older women generally have both studies, unless a cyst is likely.^{164,165,166-169} Negative imaging results place a patient a very low risk of malignancy (generally less than 3%); however, clinical judgment is necessary as some women with negative imaging may warrant biopsy and will be found to have malignant mass.^{164,170-172} The recommendations for subsequent management follow imaging assessments and clinical level of suspicion. Imaging should proceed biopsy in most situations due to potential alteration of imaging findings by the biopsy.

Symptomatic or positive findings on physical examination include palpable mass in the breast, nipple discharge without a palpable mass, asymmetric thickening or nodularity, skin changes, and breast pain.

Palpable Mass in the Breast

A palpable mass is a discrete lesion that can be readily identified during a physical exam. The NCCN Guidelines separate the evaluation of women with the palpable mass into two age groups: women aged 30 years or older and women younger than 30 years of age.

Women with Palpable Mass Aged 30 Years or Older:

The main difference in the guidelines for evaluating a palpable mass in women aged 30 years or older compared with younger women is the increased degree of suspicion of breast cancer. The initial evaluation begins with a diagnostic mammogram and ultrasound. Ultrasound should be geographically correlated with the palpable mass in question. Observation without further evaluation is not an option in these women. There are some clinical circumstances, such as mass with low clinical suspicion or suspected simple cyst, in which ultrasound would be preferred and may suffice for women 30 to 39 years of age due to the high sensitivity of ultrasound alone.^{167,168} After the diagnostic imaging assessment, the abnormality is placed into one of the following categories: negative or benign; probably benign; or suspicious or highly suggestive of cancer with management following BIRADS final assessment recommendations.

If there is a lack of geographic correlation between clinical and imaging findings, further evaluation is recommended. Sensitivity of combined mammography and ultrasound for evaluation of palpable masses is high for cancer detection, although specificity may be relatively low.

For women with mammographic findings that are suspicious or highly suggestive of breast cancer, the NCCN panel recommends ultrasound to determine lesion size and to guide tissue biopsy. The NCCN panel notes that FNA and core needle biopsy are both valuable. However, FNA requires cytologic expertise. When a core needle biopsy is utilized, concordance between the pathology report, imaging, and clinical findings must be obtained.

Ultrasound Findings:

Solid Mass:

NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

If the solid mass found on the ultrasound is suspected to be probably benign (ie, BI-RADS® category 3), the options are: 1) observation, if clinical suspicion for breast cancer is low; or 2) tissue biopsy, if the mass is clinically suspicious. Observation may be elected for those with low clinical suspicion; a physical examination follow-up with or without ultrasound or diagnostic mammogram is recommended every 6 to 12 months for 1 to 2 years to assess stability of the solid mass. There may be variability on the follow-up interval based on the level of suspicion. Numerous clinical studies now support the ability of ultrasound to accurately characterize palpable solid masses as probably benign with risk of malignancy generally less than 2%. However, these same studies have shown that many such masses will eventually warrant biopsy and compliance with follow-up may be low.^{165,167,173-177} Progression of size or suspicion on follow-up studies warrants biopsy.

Cystic Masses:

Breast cysts are either classified as simple or non-simple cysts, with the latter class being subdivided into complicated cysts and complex (cystic) mass (see Table 1 for definitions).

Simple Cyst:

A cyst meeting all criteria of a simple cyst is considered to be benign (ie, BI-RADS® 2)^{30,178} if the clinical findings and ultrasonographic results are concordant. Therapeutic fluid aspiration can be considered if clinical symptoms such as pain persist. These patients then can be followed with routine screening. Cytologic examination is recommended if bloody fluid is obtained during aspiration.

Non-Simple Complicated Cyst:

A complicated non-simple cyst is associated with a low risk of malignancy (<2%) (BI-RADS® 3).^{30,179-181} Options for managing

complicated cysts are either aspiration or short-term follow-up with physical examination and ultrasonography with or without mammography every 6 to 12 months for 1 to 2 years to assess stability. There may be variability on the follow-up interval based on the level of suspicion. The option of aspiration may be more strongly considered in a patient likely to be lost to follow-up. Complicated cysts that increase in size or suspicion should be biopsied.

If the mass resolves after aspiration, and cytology results are negative, the NCCN panel recommends that the patient should return to routine screening. If the mass first resolves after aspiration and then recurs, then repeat assessment with imaging or a surgical excision may be warranted. If the mass persists after aspiration, the NCCN panel recommends ultrasound with image-guided biopsy. Surgical excision is appropriate if unable to perform core needle biopsy.

Non-Simple Complex Cystic and Solid Mass:

A complex cystic and solid mass has both cystic and solid components. Complex cysts have a relatively high risk of malignancy (eg, 14% and 23% in 2 studies).^{30,150,180-182} The NCCN panel recommends a tissue biopsy for complex (cystic) masses (BI-RADS® 4).

No Imaging Abnormality:

If no ultrasonographic or mammographic abnormality is detected (BI-RADS® 1), tissue biopsy (core needle biopsy or excision) should be carried out for suspicious clinical findings or for low clinical suspicion and observation at 3- to 6-month intervals for 1 to 2 years should be considered to assess stability. The negative predictive value of negative imaging is high, >96%.^{164,168,171} Soo, 2001 #674,172 If the clinical lesion increases in size or suspicion, tissue sampling should be repeated, whereas routine breast screening is recommended if the lesion remains stable.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

If the option of tissue biopsy is elected, the biopsy result indicates benign mass, and this finding is concordant with the imaging results, the NCCN panel recommends a physical examination every 6 to 12 months, with or without ultrasound or mammogram, for 1 to 2 years to ensure that the lesion is stable. Routine breast screening is recommended if the lesion is stable. If the lesion increases in size, the NCCN panel recommends surgical excision.

If the diagnosis by tissue biopsy is an indeterminate lesion, atypical hyperplasia, LCIS that is non-concordant with imaging, or a benign and image discordant lesion, the NCCN panel recommends surgical excision. Mucin-producing lesions, potential phyllodes tumor, papillary lesions, radial scars, or other histologies of concern to the pathologist may also require excisional biopsy. Select patients (ie, some patients with atypical hyperplasia, LCIS, fibroepithelial lesions, radial scars) may be suitable for monitoring in lieu of surgical excision. For patients with classic LCIS that is concordant with imaging, the NCCN panel recommends routine screening along with risk reduction therapy according to the [NCCN Guidelines for Breast Cancer Risk Reduction](#) or surgical excision may be performed. Multiple-foci LCIS involving greater than 4 terminal ductal units on core biopsy is associated with increased risk of being invasive cancer.¹⁶² Patients with pleomorphic LCIS are treated with surgical excision and managed according to the [NCCN Guidelines for Breast Cancer](#).

Malignant Finding:

Malignant findings with image-guided biopsy or surgical excision should be treated according to the [NCCN Guidelines for Breast Cancer](#).

Women with Palpable Mass Younger Than 30 Years of Age:

The preferred option for initial evaluation of a palpable mass is to proceed directly to ultrasound.¹⁶⁷ Mammogram may be considered if

ultrasound or CBE results are highly suspicious or suggestive of cancer or if the patient is identified as having a high risk for breast cancer based on personal and family history. From this point, the decision tree for women younger than 30 years of age is almost identical to the pathway for older women. The main difference is consideration of a diagnostic mammogram in only some situations for the younger women. Because the incidence of malignancy in women who are younger than age 30 is low, observation of the mass for one or two menstrual cycles is also an option in cases with low clinical suspicion. If observation is elected and the mass resolves after one or two menstrual cycles, the patient may return to routine screening. If there is significant increase in size or increase in clinical suspicion, ultrasound should be performed. Needle sampling prior to imaging is not recommended.

If no ultrasonographic abnormality is found (BI-RADS® 1), a mammogram is recommended in cases where there is high clinical suspicion or for those at higher risk due to known genetic mutation or family history. Based on the mammogram results, from this point the management is identical to the pathway for older women. If the clinical suspicion is low, physical examination every 3 to 6 months for 1 to 2 years is recommended with or without ultrasound. If the mass increases in size during the observation period, diagnostic mammogram may be considered followed by tissue biopsy. If the mass remains stable, routine breast screening is recommended.

Nipple Discharge Without a Palpable Mass

Nipple discharge is common, and, in many cases, unrelated to breast pathology.¹⁸³⁻¹⁸⁹ For example, non-spontaneous discharge from multiple breast ducts in a non-lactating woman can occur during pregnancy, following breast stimulation, in women with certain thyroid conditions,



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

and in those taking certain medications, such as estrogen, oral contraceptives, opiates, and particular antihypertensive agents.¹⁸³

Suspicion of underlying pathology (eg, ductal carcinoma, papilloma) is raised when nipple discharge is persistent and reproducible on examination, spontaneous, unilateral, from a single duct, serous, sanguineous, or serosanguineous.¹⁹⁰

In patients with a nipple discharge but no palpable mass, an evaluation of the characteristics of the nipple discharge is the first step. The appropriate follow-up of a non-spontaneous, multiple-duct discharge in women younger than age 40 is observation, coupled with education of the patient to stop compression of the breast and to report the development of any spontaneous discharge. In women aged 40 years or older, mammography and a further workup based on the BI-RADS® category along with education similar to that for younger women is recommended. Evaluation of this type of nipple discharge is based on the overall BI-RADS® category of the diagnostic mammogram, if not done previously.

Women with suspicious nipple discharge are imaged with age-appropriate diagnostic mammography and ultrasound. Several clinical studies have established a very low risk of malignancy when these tests are negative.^{191,192} In certain situations, MRI or ductography may play an adjunctive role, aiding in identifying a possible abnormality and its location. Several studies have shown that breast MRI aids in the diagnosis of suspected ductal disease.^{135-137,193,194}

According to the NCCN panel, when an overall imaging BI-RADS® assessment is category 1-3, either a ductogram or MRI are optional to guide the duct excision. The management options include duct excision¹⁹⁵ or follow-up with physical exam after 6 months and imaging

with diagnostic mammogram with or without ultrasound for 1 to 2 years. If clinical suspicion increases during follow-up, tissue biopsy is recommended.

For BI-RADS® category 4 or 5, the NCCN panel recommends a tissue biopsy. If the biopsy findings are indeterminate, a ductogram is optional, but surgical duct excision would still be necessary. If findings are indicative of malignancy, the patient should be treated according to the [NCCN Guidelines for Breast Cancer](#).

Asymmetric Thickening or Nodularity

Thickening, nodularity, or asymmetry is distinct from a palpable mass in that the finding is ill-defined and often vague on physical breast examination. Factors to consider include whether the thickening is a new or previous finding, and whether or not it appears to be representative of normal asymmetry. Imaging evaluation follows that of a palpable mass.¹⁶⁴ If the patient is younger than age 30 years and has no high risk factors, ultrasound evaluation is appropriate followed by consideration of diagnostic mammography. Diagnostic mammograms for this age group are fairly low in yield because of the density of the breast and low risk of breast cancer. In a woman aged 30 years or older, a diagnostic mammogram and an ultrasound evaluation should be obtained.

If the overall imaging findings are classified as BI-RADS® category 1-3 *and* the clinical assessment is benign, the patient should be clinically reexamined with imaging as needed in 3 to 6 months to assess stability. Age-appropriate diagnostic mammogram and/or ultrasound may be performed every 6 to 12 months for 1 to 2 years to assess stability. If the findings on physical exam and/or imaging are stable, routine screening can be resumed. If the either or both findings indicate



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

progression, it should be investigated as previously described for palpable mass.

If a clinically suspicious change is noted or the overall imaging findings are classified as BI-RADS® assessment category 4-5, a tissue biopsy is recommended.

Skin Changes

Any type of unusual skin changes around the breast may represent serious disease and needs evaluation. IBC should be considered when dermal edema (peau d'orange) and breast erythema are present, and nipple excoriation, scaling, and eczema should increase clinical suspicion of Paget's disease. IBC is a rare, aggressive form of breast cancer estimated to account for 1% to 6% of breast cancer cases in the United States. IBC is a clinical diagnosis that requires erythema and dermal edema of a third or more of the skin of the breast with a palpable border to the erythema.^{196,197} Paget's disease of the breast is a rare manifestation of breast cancer characterized by neoplastic cells in the epidermis of the nipple areolar complex. It most commonly presents with eczema of the nipple or areola, bleeding, ulceration, and itching of the nipple. The diagnosis is often delayed because of the rare nature of the condition and confusion with other dermatologic conditions.^{198,199} Pure Paget's disease is frequently occult on mammography²⁰⁰ and a negative mammogram does not exclude Paget's disease, which requires skin biopsy.

The initial evaluation of a patient with breast skin changes begins with a bilateral diagnostic mammogram with or without ultrasound imaging. If the imaging results are abnormal, the evaluation proceeds on the basis of the imaging findings. If the breast imaging results are normal, further workup is still needed.

Punch biopsy of the skin or nipple biopsy should be performed following imaging findings consistent with an overall BI-RADS® assessment category 1-3. Antibiotics may or may not be given, depending on the clinical suspicion for breast infection, but should not delay diagnostic evaluation. If biopsy results are benign, clinical and pathological correlation should be reassessed. In addition, a breast MRI, a repeat biopsy, and consultation with a breast specialist should be considered. If the skin biopsy is malignant, the patient should be treated according to the [NCCN Guidelines for Breast Cancer](#).

A tissue biopsy should be performed if imaging findings are consistent of an overall BI-RADS® assessment category 4-5. According to the NCCN panel, core needle biopsy is the preferred option with or without punch biopsy, although surgical excision is also an option. A benign biopsy result should be followed by a punch biopsy of the skin, if not previously performed, or nipple biopsy, with reassessment as described above for BI-RADS® category 1-3. A biopsy showing a malignant finding should be managed according to the [NCCN Guidelines for Breast Cancer](#).

Breast Pain

Breast pain is the most common symptom in the breast. Individuals presenting with breast pain fear that this is a symptom of breast cancer, therefore causing significant anxiety. The NCCN panel has developed guidelines to evaluate breast pain for cancer and provide reassurance. The risk of cancer in a woman presenting with breast pain as the only symptom is low, between 1.2% and 6.7%.^{4,123,201,202}

Evaluation of breast pain includes comprehensive history, type of pain, relationship to menses, duration, location, impact on activities of daily living, factors that aggravate/alleviate pain, any other medical problems



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

and comorbidities, and a thorough CBE. If CBE fails to identify any physical abnormality such as palpable mass, asymmetric thickening, nipple discharge, or skin changes; the pain is cyclic; or diffuse and non-focal and screening mammograms are current and negative, the NCCN panel recommends providing reassurance to the patient and treating the pain with symptomatic management (eg, over-the-counter pain medications, if needed; use of a good support bra; ice packs or heating pads). Cyclical breast pain may often spontaneously resolve. Reassurance alone has shown to help resolve the symptom in 86% of women with mild pain and in 52% of women with severe pain.²⁰³

If the breast pain is focal and persistent in nature, the NCCN panel recommends age-appropriate diagnostic imaging (diagnostic mammogram with or without ultrasound for those ≥ 30 years of age; and ultrasound for those < 30 years of age).

For those with BI-RADS® assessment category 1, the panel recommends a clinical encounter every 6 to 12 months for 1 to 2 years along with symptomatic management of the breast pain, if desired. For a simple cyst (BI-RADS® assessment category 2) geographically correlated with focal pain, drainage may be considered for symptom relief. Aspiration is recommended for painful, complicated cysts (BI-RADS 3) to rule out infection or malignancy. A tissue biopsy should be performed if imaging findings are consistent of an overall BI-RADS® assessment category 4-5.

Summary

The intent of the NCCN Breast Cancer Screening and Diagnosis guidelines is to give health care providers a practical, consistent framework for screening and evaluating a spectrum of clinical breast lesions. Clinical judgment should always be an important component of the optimal management of the patient.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

Table 1: Breast Cysts - Types and Definitions

Simple cyst	Anechoic (cystic), well-circumscribed, round or oval with well-defined imperceptible wall and posterior enhancement.
Non-simple cyst	Has one or more characteristics not found in a simple cyst.
<ul style="list-style-type: none"> • complicated 	Has most but not all elements of a simple cyst. Complicated cysts do not contain solid elements, intracystic masses, thick walls, or thick septa. This type of cyst may contain low-level echoes or intracystic debris, and can be described as a round, circumscribed mass containing low-level echoes without vascular flow, fulfilling most but not all criteria of a simple cyst.
<ul style="list-style-type: none"> • complex 	Has some discrete solid component, which may include thick walls, thick septa, and/or intracystic mass. Complex cysts have both anechoic (cystic) and echogenic (solid) components.
References	139,150,178-182,204

Discussion
Update in
progress



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

References

1. American Cancer Society. Breast Cancer Facts and Figures 2009-2010. Atlanta: American Cancer Society, Inc. Available at: <http://www.cancer.org/Research/CancerFactsFigures/BreastCancerFactsFigures/breast-cancer-facts--figures-2009-2010>.
2. Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. CA Cancer J Clin 2017;67:7-30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28055103>.
3. Humphrey LL, Helfand M, Chan BK, Woolf SH. Breast cancer screening: a summary of the evidence for the U.S. Preventive Services Task Force. Ann Intern Med 2002;137:347-360. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12204020>.
4. Barton MB, Harris R, Fletcher SW. The rational clinical examination. Does this patient have breast cancer? The screening clinical breast examination: should it be done? How? JAMA 1999;282:1270-1280. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10517431>.
5. Bevers TB. Breast awareness: a shift in the paradigm of breast self-examination. J Natl Compr Canc Netw 2009;7:1042-1043. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1993097>.
6. Bevers T. Breast Self-examination. In: Singletary SE, Robb GL, Hortobagyi GN, eds. Advanced Therapy of Breast Disease. 2nd ed. New York: B.C. Decker, Inc; 2004.
7. Thomas DB, Gao DL, Ray RM, et al. Randomized trial of breast self-examination in Shanghai: final results. J Natl Cancer Inst 2002;94:1445-1457. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12359854>.
8. Feig SA, Yaffe MJ. Digital mammography. Radiographics 1998;18:893-901. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9672974>.
9. Pisano ED, Yaffe MJ, Hemminger BM, et al. Current status of full-field digital mammography. Acad Radiol 2000;7:266-280. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10766101>.
10. Pisano ED, Gatsonis C, Hendrick E, et al. Diagnostic performance of digital versus film mammography for breast-cancer screening. N Engl J Med 2005;353:1773-1783. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16169887>.
11. Pisano ED, Hendrick RE, Yaffe MJ, et al. Diagnostic accuracy of digital versus film mammography: exploratory analysis of selected population subgroups in DMIST. Radiology 2008;246:376-383. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18227537>.
12. Skaane P, Hofvind S, Skjennald A. Randomized trial of screen-film versus full-field digital mammography with soft-copy reading in population-based screening program: follow-up and final results of Oslo II study. Radiology 2007;244:708-717. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17709826>.
13. Ciatto S, Houssami N, Bernardi D, et al. Integration of 3D digital mammography with tomosynthesis for population breast-cancer screening (STORM): a prospective comparison study. Lancet Oncol 2013;14:583-589. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23623721>.
14. Skaane P, Bandos AI, Gullien R, et al. Comparison of digital mammography alone and digital mammography plus tomosynthesis in a population-based screening program. Radiology 2013;267:47-56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23297332>.
15. Rafferty EA, Park JM, Philpotts LE, et al. Assessing radiologist performance using combined digital mammography and breast tomosynthesis compared with digital mammography alone: results of a multicenter, multireader trial. Radiology 2013;266:104-113. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23169790>.
16. Friedewald SM, Rafferty EA, Rose SL, et al. Breast cancer screening using tomosynthesis in combination with digital mammography. JAMA 2014;311:2499-2507. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25058084>.

17. Lourenco AP, Barry-Brooks M, Baird GL, et al. Changes in recall type and patient treatment following implementation of screening digital breast tomosynthesis. *Radiology* 2015;274:337-342. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25247407>.
18. Rose SL, Tidwell AL, Ice MF, et al. A reader study comparing prospective tomosynthesis interpretations with retrospective readings of the corresponding FFDM examinations. *Acad Radiol* 2014;21:1204-1210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25107868>.
19. Destounis S, Arieno A, Morgan R. Initial experience with combination digital breast tomosynthesis plus full field digital mammography or full field digital mammography alone in the screening environment. *J Clin Imaging Sci* 2014;4:9. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24744966>.
20. Margolies L, Cohen A, Sonnenblick E, et al. Digital breast tomosynthesis changes management in patients seen at a tertiary care breast center. *ISRN Radiol* 2014;2014:658929. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24967297>.
21. Lang K, Andersson I, Rosso A, et al. Performance of one-view breast tomosynthesis as a stand-alone breast cancer screening modality: results from the Malmö Breast Tomosynthesis Screening Trial, a population-based study. *Eur Radiol* 2016;26:184-190. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25929946>.
22. Gilbert FJ, Tucker L, Gillan MG, et al. Accuracy of digital breast tomosynthesis for depicting breast cancer subgroups in a UK retrospective reading study (TOMMY Trial). *Radiology* 2015;277:697-706. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26176654>.
23. Zuckerman SP, Conant EF, Keller BM, et al. Implementation of synthesized two-dimensional mammography in a population-based digital breast tomosynthesis screening program. *Radiology* 2016;160366. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27467468>.
24. Skaane P, Bandos AI, Eben EB, et al. Two-view digital breast tomosynthesis screening with synthetically reconstructed projection images: comparison with digital breast tomosynthesis with full-field digital mammographic images. *Radiology* 2014;271:655-663. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24484063>.
25. Nelson HD, Zakher B, Cantor A, et al. Risk factors for breast cancer for women aged 40 to 49 years: a systematic review and meta-analysis. *Ann Intern Med* 2012;156:635-648. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22547473>.
26. Mandelson MT, Oestreicher N, Porter PL, et al. Breast density as a predictor of mammographic detection: comparison of interval- and screen-detected cancers. *J Natl Cancer Inst* 2000;92:1081-1087. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10880551>.
27. Chiu SY, Duffy S, Yen AM, et al. Effect of baseline breast density on breast cancer incidence, stage, mortality, and screening parameters: 25-year follow-up of a Swedish mammographic screening. *Cancer Epidemiol Biomarkers Prev* 2010;19:1219-1228. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20406961>.
28. Vachon CM, Sellers TA, Scott CG, et al. Longitudinal breast density and risk of breast cancer [abstract]. *Cancer Research* 2011;70:Abstract 4828. Available at: http://cancerres.aacrjournals.org/cgi/content/meeting_abstract/70/8_MeetingAbstracts/4828.
29. Richman I, Asch SM, Bendavid E, et al. Breast density notification legislation and breast cancer stage at diagnosis: early evidence from the SEER registry. *J Gen Intern Med* 2016. Available at:
30. Berg WA, Blume JD, Cormack JB, et al. Combined screening with ultrasound and mammography vs mammography alone in women at elevated risk of breast cancer. *JAMA* 2008;299:2151-2163. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18477782>.
31. Berg WA, Zhang Z, Lehrer D, et al. Detection of breast cancer with addition of annual screening ultrasound or a single screening MRI to



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

mammography in women with elevated breast cancer risk. JAMA 2012;307:1394-1404. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22474203>.

32. Berg WA, Bandos AI, Mendelson EB, et al. Ultrasound as the Primary Screening Test for Breast Cancer: Analysis From ACRIN 6666. J Natl Cancer Inst 2016;108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26712110>.

33. Weigert J, Steenbergen S. The connecticut experiments second year: ultrasound in the screening of women with dense breasts. Breast J 2015;21:175-180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25683369>.

34. Bevers TB. Ultrasound for the screening of breast cancer. Curr Oncol Rep 2008;10:527-528. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18928668>.

35. Hooley RJ, Greenberg KL, Stackhouse RM, et al. Screening US in patients with mammographically dense breasts: initial experience with Connecticut Public Act 09-41. Radiology 2012;265:59-69. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22723501>.

36. Scheel JR, Lee JM, Sprague BL, et al. Screening ultrasound as an adjunct to mammography in women with mammographically dense breasts. Am J Obstet Gynecol 2015;212:9-17. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24959654>.

37. Lord SJ, Lei W, Craft P, et al. A systematic review of the effectiveness of magnetic resonance imaging (MRI) as an addition to mammography and ultrasound in screening young women at high risk of breast cancer. Eur J Cancer 2007;43:1905-1917. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17681781>.

38. Mann RM, Kuhl CK, Kinkel K, Boetes C. Breast MRI: guidelines from the European Society of Breast Imaging. Eur Radiol 2008;18:1307-1318. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18389253>.

39. Schnall M, Orel S. Breast MR imaging in the diagnostic setting. Magn Reson Imaging Clin N Am 2006;14:329-337, vi. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17098174>.

40. Ng AK, Garber JE, Diller LR, et al. Prospective study of the efficacy of breast magnetic resonance imaging and mammographic screening in survivors of hodgkin lymphoma. J Clin Oncol 2013;31:2282-2288. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23610104>.

41. Kuhl CK, Schrading S, Leutner CC, et al. Mammography, breast ultrasound, and magnetic resonance imaging for surveillance of women at high familial risk for breast cancer. J Clin Oncol 2005;23:8469-8476. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16293877>.

42. Kriege M, Brekelmans CTM, Boetes C, et al. Differences between first and subsequent rounds of the MRISC breast cancer screening program for women with a familial or genetic predisposition. Cancer 2006;106:2318-2326. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16615112>.

43. Kriege M, Brekelmans CTM, Obdeijn IM, et al. Factors affecting sensitivity and specificity of screening mammography and MRI in women with an inherited risk for breast cancer. Breast Cancer Res Treat 2006;100:109-119. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16791481>.

44. Robson ME, Offit K. Breast MRI for women with hereditary cancer risk. JAMA 2004;292:1368-1370. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15367560>.

45. Warner E. The role of magnetic resonance imaging in screening women at high risk of breast cancer. Top Magn Reson Imaging 2008;19:163-169. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18941396>.

46. Warner E, Messersmith H, Causer P, et al. Systematic review: using magnetic resonance imaging to screen women at high risk for breast cancer. Ann Intern Med 2008;148:671-679. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18458280>.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

47. Warner E, Plewes DB, Hill KA, et al. Surveillance of BRCA1 and BRCA2 mutation carriers with magnetic resonance imaging, ultrasound, mammography, and clinical breast examination. JAMA 2004;292:1317-1325. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15367553>.

48. Lehman CD, Smith RA. The role of MRI in breast cancer screening. J Natl Compr Canc Netw 2009;7:1109-1115. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19930977>.

49. Saslow D, Boetes C, Burke W, et al. American Cancer Society guidelines for breast screening with MRI as an adjunct to mammography. CA Cancer J Clin 2007;57:75-89. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17392385>.

50. Leach MO, Boggis CR, Dixon AK, et al. Screening with magnetic resonance imaging and mammography of a UK population at high familial risk of breast cancer: a prospective multicentre cohort study (MARIBS). Lancet 2005;365:1769-1778. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15910949>.

51. Port ER, Park A, Borgen PI, et al. Results of MRI screening for breast cancer in high-risk patients with LCIS and atypical hyperplasia. Ann Surg Oncol 2007;14:1051-1057. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17206485>.

52. Ramalho J, Ramalho M, Jay M, et al. Gadolinium Toxicity and Treatment. Magn Reson Imaging 2016. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27693607>.

53. Stojanov D, Aracki-Trenkic A, Benedeto-Stojanov D. Gadolinium deposition within the dentate nucleus and globus pallidus after repeated administrations of gadolinium-based contrast agents-current status. Neuroradiology 2016;58:433-441. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26873830>.

54. Xia D, Davis RL, Crawford JA, Abraham JL. Gadolinium released from MR contrast agents is deposited in brain tumors: in situ demonstration using scanning electron microscopy with energy

dispersive X-ray spectroscopy. Acta Radiol 2010;51:1126-1136. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20868305>.

55. ACR practise guideline for the performance of contrast-enhanced magnetic resonance imaging (MRI) of the breast 2012. Available at: <http://www.acr.org/Search?q=guidelines%20for%20contrast-enhanced%20MRI%20of%20the%20breast>.

56. Hruska CB. Molecular breast imaging for screening in dense breasts: state of the art and future directions. AJR Am J Roentgenol 2017;208:275-283. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27762607>.

57. Shermis RB, Wilson KD, Doyle MT, et al. Supplemental breast cancer screening with molecular breast imaging for women with dense breast tissue. AJR Am J Roentgenol 2016;207:450-457. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27186635>.

58. Meissnitzer T, Seymer A, Keinrath P, et al. Added value of semi-quantitative breast-specific gamma imaging in the work-up of suspicious breast lesions compared to mammography, ultrasound and 3-T MRI. Br J Radiol 2015;88:20150147. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25882690>.

59. Jochelson M. Contrast-enhanced digital mammography. Radiol Clin North Am 2014;52:609-616. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24792660>.

60. U.S. Food and Drug Administration (FDA) safety alert on ductal lavage. Available at: <http://www.breastcancer.org/research-news/20140220>

61. Oeffinger KC, Fontham ET, Etzioni R, et al. Breast Cancer Screening for Women at Average Risk: 2015 Guideline Update From the American Cancer Society. JAMA 2015;314:1599-1614. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26501536>.

62. Siu AL, Force USPST. Screening for Breast Cancer: U.S. Preventive Services Task Force Recommendation Statement. Ann

Intern Med 2016;164:279-296. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26757170>.

63. Myers ER, Moorman P, Gierisch JM, et al. Benefits and harms of breast cancer screening: A systematic review. JAMA 2015;314:1615-1634. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26501537>.

64. Moss SM, Cuckle H, Evans A, et al. Effect of mammographic screening from age 40 years on breast cancer mortality at 10 years' follow-up: a randomised controlled trial. Lancet 2006;368:2053-2060. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17161727>.

65. Moss SM, Wale C, Smith R, et al. Effect of mammographic screening from age 40 years on breast cancer mortality in the UK Age trial at 17 years' follow-up: a randomised controlled trial. Lancet Oncol 2015;16:1123-1132. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/26206144>.

66. Hellquist BN, Duffy SW, Abdsaleh S, et al. Effectiveness of population-based service screening with mammography for women ages 40 to 49 years: evaluation of the Swedish Mammography Screening in Young Women (SCRY) cohort. Cancer 2011;117:714-722. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20882563>.

67. Swedish Organised Service Screening Evaluation G. Reduction in breast cancer mortality from organized service screening with mammography: 1. Further confirmation with extended data. Cancer Epidemiol Biomarkers Prev 2006;15:45-51. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/16434585>.

68. Coldman A, Phillips N, Wilson C, et al. Pan-Canadian study of mammography screening and mortality from breast cancer. J Natl Cancer Inst 2014;106. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25274578>.

69. Broeders M, Moss S, Nystrom L, et al. The impact of mammographic screening on breast cancer mortality in Europe: a review of observational studies. J Med Screen 2012;19 Suppl 1:14-25. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22972807>.

70. Malmgren JA, Parikh J, Atwood MK, Kaplan HG. Impact of mammography detection on the course of breast cancer in women aged 40-49 years. Radiology 2012;262:797-806. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22357883>.

71. Saadatmand S, Bretveld R, Siesling S, Tilanus-Linthorst MM. Influence of tumour stage at breast cancer detection on survival in modern times: population based study in 173,797 patients. BMJ 2015;351:h4901. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26442924>.

72. Mandelblatt JS, Cronin KA, Bailey S, et al. Effects of mammography screening under different screening schedules: model estimates of potential benefits and harms. Ann Intern Med 2009;151:738-747. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19920274>.

73. Modeling Report: Collaborative Modeling of U.S. Breast Cancer Screening Strategies: Breast Cancer: Screening. U.S. Preventive Services Task Force. April 2015. Available at:
<http://www.uspreventiveservicestaskforce.org/Page/Document/modeling-report-collaborative-modeling-of-us-breast-cancer-1/breast-cancer-screening1>.

74. American College of O-G. Practice bulletin no. 122: Breast cancer screening. Obstet Gynecol 2011;118:372-382. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/21775869>.

75. Schwartz LM, Woloshin S, Sox HC, et al. US women's attitudes to false positive mammography results and detection of ductal carcinoma in situ: cross sectional survey. BMJ 2000;320:1635-1640. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/10856064>.

76. Coldman AJ, Phillips N, Speers C. A retrospective study of the effect of participation in screening mammography on the use of chemotherapy and breast conserving surgery. Int J Cancer 2007;120:2185-2190. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17290404>.

77. Tosteson AN, Fryback DG, Hammond CS, et al. Consequences of false-positive screening mammograms. JAMA Intern Med 2014;174:954-961. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24756610>.

78. Yen AM, Duffy SW, Chen TH, et al. Long-term incidence of breast cancer by trial arm in one county of the Swedish Two-County Trial of mammographic screening. Cancer 2012;118:5728-5732. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22605639>.

79. Bleyer A, Welch HG. Effect of three decades of screening mammography on breast-cancer incidence. N Engl J Med 2012;367:1998-2005. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23171096>.

80. Puliti D, Duffy SW, Miccinesi G, et al. Overdiagnosis in mammographic screening for breast cancer in Europe: a literature review. J Med Screen 2012;19 Suppl 1:42-56. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22972810>.

81. Zackrisson S, Andersson I, Janzon L, et al. Rate of over-diagnosis of breast cancer 15 years after end of Malmö mammographic screening trial: follow-up study. BMJ 2006;332:689-692. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/16517548>.

82. Miller AB, To T, Baines CJ, Wall C. Canadian National Breast Screening Study-2: 13-year results of a randomized trial in women aged 50-59 years. J Natl Cancer Inst 2000;92:1490-1499. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/10995804>.

83. Miller AB, To T, Baines CJ, Wall C. The Canadian National Breast Screening Study-1: breast cancer mortality after 11 to 16 years of follow-up. A randomized screening trial of mammography in women age 40 to 49 years. Ann Intern Med 2002;137:305-312. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/12204013>.

84. Marmot MG, Altman DG, Cameron DA, et al. The benefits and harms of breast cancer screening: an independent review. Br J Cancer

2013;108:2205-2240. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23744281>.

85. Gunsoy NB, Garcia-Closas M, Moss SM. Estimating breast cancer mortality reduction and overdiagnosis due to screening for different strategies in the United Kingdom. Br J Cancer 2014;110:2412-2419. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24762956>.

86. Johns LE, Coleman DA, Swerdlow AJ, Moss SM. Effect of population breast screening on breast cancer mortality up to 2005 in England and Wales: an individual-level cohort study. Br J Cancer 2017;116:246-252. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27931047>.

87. Mandelblatt JS, Stout NK, Schechter CB, et al. Collaborative modeling of the benefits and harms associated with different U.S. breast cancer screening strategies. Ann Intern Med 2016;164:215-225. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26756606>.

88. Hendrick RE, Helvie MA, Hardesty LA. Implications of CISNET modeling on number needed to screen and mortality reduction with digital mammography in women 40-49 years old. AJR Am J Roentgenol 2014;203:1379-1381. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25415718>.

89. Mandelblatt, JS.; Cronin, K.; de Koning, H.; Miglioretti, DL.; Schechter, CS.; Stout, N. Collaborative modeling of U.S. breast cancer screening strategies. AHRQ Publication No. 14-05201-EF-4. 2015. Available at:

<https://www.uspreventiveservicestaskforce.org/Page/Document/modeling-report-collaborative-modeling-of-us-breast-cancer-1/breast-cancer-screening1>.

90. Engel JM, Stankowski-Drengler TJ, Stankowski RV, et al. All-cause mortality is decreased in women undergoing annual mammography before breast cancer diagnosis. AJR Am J Roentgenol 2015;204:898-902. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25794084>.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

91. Duffy SW, Diben A, Michalopoulos D, et al. Screen detection of ductal carcinoma in situ and subsequent incidence of invasive interval breast cancers: a retrospective population-based study. *Lancet Oncol* 2016;17:109-114. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26655422>.

92. Badgwell BD, Giordano SH, Duan ZZ, et al. Mammography before diagnosis among women age 80 years and older with breast cancer. *J Clin Oncol* 2008;26:2482-2488. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18427152>.

93. Mandelblatt JS, Silliman R. Hanging in the balance: making decisions about the benefits and harms of breast cancer screening among the oldest old without a safety net of scientific evidence. *J Clin Oncol* 2009;27:487-490. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19075258>.

94. van Dijck J, Verbeek A, Hendriks J, et al. Mammographic screening after the age of 65 years: early outcomes in the Nijmegen programme. *Br J Cancer* 1996;74:1838-1842. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8956803>.

95. Walter LC, Covinsky KE. Cancer screening in elderly patients: a framework for individualized decision making. *JAMA* 2001;285:2750-2756. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11386931>.

96. Smith RA, Saslow D, Sawyer KA, et al. American Cancer Society guidelines for breast cancer screening: update 2003. *CA Cancer J Clin* 2003;53:141-169. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12809408>.

97. Costantino JP, Gail MH, Pee D, et al. Validation studies for models projecting the risk of invasive and total breast cancer incidence. *J Natl Cancer Inst* 1999;91:1541-1548. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10491430>.

98. Gail MH, Brinton LA, Byar DP, et al. Projecting individualized probabilities of developing breast cancer for white females who are

being examined annually. *J Natl Cancer Inst* 1989;81:1879-1886. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2593165>.

99. Gail MH, Costantino JP. Validating and improving models for projecting the absolute risk of breast cancer. *J Natl Cancer Inst* 2001;93:334-335. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11238688>.

100. Rockhill B, Spiegelman D, Byrne C, et al. Validation of the Gail et al. model of breast cancer risk prediction and implications for chemoprevention. *J Natl Cancer Inst* 2001;93:358-366. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11238697>.

101. Spiegelman D, Colditz GA, Hunter D, Hertzmark E. Validation of the Gail et al. model for predicting individual breast cancer risk. *J Natl Cancer Inst* 1994;86:600-607. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8145275>.

102. Gail MH, Costantino JP, Pee D, et al. Projecting individualized absolute invasive breast cancer risk in African American women. *J Natl Cancer Inst* 2007;99:1782-1792. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18042936>.

103. Matsuno RK, Costantino JP, Ziegler RG, et al. Projecting individualized absolute invasive breast cancer risk in Asian and Pacific Islander American women. *J Natl Cancer Inst* 2011. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21562243>.

104. Arpino G, Laucirica R, Elledge RM. Premalignant and in situ breast disease: biology and clinical implications. *Ann Intern Med* 2005;143:446-457. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16172443>.

105. Lakhani SR. In-situ lobular neoplasia: time for an awakening. *Lancet* 2003;361:96. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12531570>.

106. Page DL, Schuyler PA, Dupont WD, et al. Atypical lobular hyperplasia as a unilateral predictor of breast cancer risk: a



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

retrospective cohort study. Lancet 2003;361:125-129. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12531579>.

107. Claus EB, Risch N, Thompson WD. Autosomal dominant inheritance of early-onset breast cancer. Implications for risk prediction. Cancer 1994;73:643-651. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8299086>.

108. Tyrer J, Duffy SW, Cuzick J. A breast cancer prediction model incorporating familial and personal risk factors. Stat Med 2004;23:1111-1130. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15057881>.

109. Amir E, Evans DG, Shenton A, et al. Evaluation of breast cancer risk assessment packages in the family history evaluation and screening programme. J Med Genet 2003;40:807-814. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14627668>.

110. Evans DG, Howell A. Breast cancer risk-assessment models. Breast Cancer Res 2007;9:213. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17888188>.

111. Saslow D, Boetes C, Burke W, et al. American Cancer Society guidelines for breast screening with MRI as an adjunct to mammography. CA Cancer J Clin 2007;57:75-89. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17392385>.

112. Parmigiani G, Berry D, Aguilar O. Determining carrier probabilities for breast cancer-susceptibility genes BRCA1 and BRCA2. Am J Hum Genet 1998;62:145-158. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9443863>.

113. Antoniou AC, Cunningham AP, Peto J, et al. The BOADICEA model of genetic susceptibility to breast and ovarian cancers: updates and extensions. Br J Cancer 2008;98:1457-1466. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18349832>.

114. Bhatia S, Robison LL, Oberlin O, et al. Breast cancer and other second neoplasms after childhood Hodgkin's disease. N Engl J Med

1996;334:745-751. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8592547>.

115. Bhatia S, Yasui Y, Robison LL, et al. High risk of subsequent neoplasms continues with extended follow-up of childhood Hodgkin's disease: report from the Late Effects Study Group. J Clin Oncol 2003;21:4386-4394. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14645429>.

116. Hancock SL, Tucker MA, Hoppe RT. Breast cancer after treatment of Hodgkin's disease. J Natl Cancer Inst 1993;85:25-31. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8416252>.

117. Metayer C, Lynch CF, Clarke EA, et al. Second cancers among long-term survivors of Hodgkin's disease diagnosed in childhood and adolescence. J Clin Oncol 2000;18:2435-2443. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10856104>.

118. van Leeuwen FE, Klokman WJ, Stovall M, et al. Roles of radiation dose, chemotherapy, and hormonal factors in breast cancer following Hodgkin's disease. J Natl Cancer Inst 2003;95:971-980. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12837833>.

119. Yahalom J, Petrek JA, Biddinger PW, et al. Breast cancer in patients irradiated for Hodgkin's disease: a clinical and pathologic analysis of 45 events in 37 patients. J Clin Oncol 1992;10:1674-1681. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1403050>.

120. Travis LB, Hill D, Dores GM, et al. Cumulative absolute breast cancer risk for young women treated for Hodgkin lymphoma. J Natl Cancer Inst 2005;97:1428-1437. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16204692>.

121. Oeffinger KC, Ford JS, Moskowitz CS, et al. Breast cancer surveillance practices among women previously treated with chest radiation for a childhood cancer. JAMA 2009;301:404-414. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19176442>.



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

122. Children's Oncology Group Long-term Follow-up Guidelines. Version 3. 2008. Available at:

123. Barton MB, Elmore JG, Fletcher SW. Breast symptoms among women enrolled in a health maintenance organization: frequency, evaluation, and outcome. *Ann Intern Med* 1999;130:651-657. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10215561>.

124. Noroozian M, Hadjiiski L, Rahnama-Moghadam S, et al. Digital breast tomosynthesis is comparable to mammographic spot views for mass characterization. *Radiology* 2012;262:61-68. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21998048>.

125. Zuley ML, Bandos AI, Ganott MA, et al. Digital breast tomosynthesis versus supplemental diagnostic mammographic views for evaluation of noncalcified breast lesions. *Radiology* 2013;266:89-95. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23143023>.

126. Waldherr C, Cerny P, Altermatt HJ, et al. Value of one-view breast tomosynthesis versus two-view mammography in diagnostic workup of women with clinical signs and symptoms and in women recalled from screening. *AJR Am J Roentgenol* 2013;200:226-231. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23255766>.

127. Food and Drug Administration. Quality mammography standards; correction—FDA: final rule; correction. 62 Federal Register 60613-60632 (1997). Available at:

128. American College of Radiology (ACR). ACR-BI-RADS®--5th Edition. ACR Breast Imaging Reporting and Data System, Breast Imaging Atlas; BI-RADS. Reston VA. American College of Radiology, 2014. . Available at: www.acr.org.

129. Bassett L, Winchester DP, Caplan RB, et al. Stereotactic core-needle biopsy of the breast: a report of the Joint Task Force of the American College of Radiology, American College of Surgeons, and College of American Pathologists. *CA Cancer J Clin* 1997;47:171-190. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9152175>.

130. Flobbe K, Bosch AM, Kessels AG, et al. The additional diagnostic value of ultrasonography in the diagnosis of breast cancer. *Arch Intern Med* 2003;163:1194-1199. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/12767956>.

131. Buchberger W, DeKoekoek-Doll P, Springer P, et al. Incidental findings on sonography of the breast: clinical significance and diagnostic workup. *AJR Am J Roentgenol* 1999;173:921-927. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10511149>.

132. Corsetti V, Houssami N, Ferrari A, et al. Breast screening with ultrasound in women with mammography-negative dense breasts: evidence on incremental cancer detection and false positives, and associated cost. *Eur J Cancer* 2008;44:539-544. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18267357>.

133. Kaplan SS. Clinical utility of bilateral whole-breast US in the evaluation of women with dense breast tissue. *Radiology* 2001;221:641-649. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11719658>.

134. Renz DM, Baltzer PAT, Bottcher J, et al. Inflammatory breast carcinoma in magnetic resonance imaging: a comparison with locally advanced breast cancer. *Acad Radiol* 2008;15:209-221. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18206620>.

135. Bahl M, Baker JA, Greenup RA, Ghate SV. Evaluation of pathologic nipple discharge: What is the added diagnostic value of MRI? *Ann Surg Oncol* 2015;22 Suppl 3:S435-441. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26249144>.

136. Lubina N, Schedelbeck U, Roth A, et al. 3.0 Tesla breast magnetic resonance imaging in patients with nipple discharge when mammography and ultrasound fail. *Eur Radiol* 2015;25:1285-1293. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25433415>.

137. Morrogh M, Morris EA, Liberman L, et al. MRI identifies otherwise occult disease in select patients with Paget disease of the nipple. *J Am*



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

Coll Surg 2008;206:316-321. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/18222386>.

138. ., Abati A, Simsir A. Breast fine needle aspiration biopsy: prevailing recommendations and contemporary practices. Clin Lab Med 2005;25:631-654. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16308084>.

139. Levine P, Simsir A, Cangiarella J. Management issues in breast lesions diagnosed by fine-needle aspiration and percutaneous core breast biopsy. Am J Clin Pathol 2006;125 Suppl:S124-134. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16830962>.

140. Pijnappel RM, van den Donk M, Holland R, et al. Diagnostic accuracy for different strategies of image-guided breast intervention in cases of nonpalpable breast lesions. Br J Cancer 2004;90:595-600. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14760370>.

141. Pisano ED, Fajardo LL, Caudry DJ, et al. Fine-needle aspiration biopsy of nonpalpable breast lesions in a multicenter clinical trial: results from the radiologic diagnostic oncology group V. Radiology 2001;219:785-792. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11376270>.

142. Fishman JE, Milikowski C, Ramsinghani R, et al. US-guided core-needle biopsy of the breast: how many specimens are necessary? Radiology 2003;226:779-782. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12601206>.

143. Yeow KM, Lo YF, Wang CS, et al. Ultrasound-guided core needle biopsy as an initial diagnostic test for palpable breast masses. J Vasc Interv Radiol 2001;12:1313-1317. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11698631>.

144. Verkooijen HM. Diagnostic accuracy of stereotactic large-core needle biopsy for nonpalpable breast disease: results of a multicenter prospective study with 95% surgical confirmation. Int J Cancer 2002;99:853-859. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12115488>.

145. Pandelidis S, Heiland D, Jones D, et al. Accuracy of 11-gauge vacuum-assisted core biopsy of mammographic breast lesions. Ann Surg Oncol 2003;10:43-47. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12513959>.

146. Kettritz U, Rotter K, Schreer I, et al. Stereotactic vacuum-assisted breast biopsy in 2874 patients: a multicenter study. Cancer 2004;100:245-251. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/14716757>.

147. Sigal-Zafrani B, Muller K, El Khoury C, et al. Vacuum-assisted large-core needle biopsy (VLNB) improves the management of patients with breast microcalcifications - analysis of 1009 cases. Eur J Surg Oncol 2008;34:377-381. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17604937>.

148. Burbank F, Forcier N. Tissue marking clip for stereotactic breast biopsy: initial placement accuracy, long-term stability, and usefulness as a guide for wire localization. Radiology 1997;205:407-415. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/9356621>.

149. Egyed Z, Pentek Z, Jaray B, et al. Radial scar-significant diagnostic challenge. Pathol Oncol Res 2008;14:123-129. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18409019>.

150. Tea M-KM, Grimm C, Fink-Retter A, et al. The validity of complex breast cysts after surgery. Am J Surg 2009;197:199-202. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18649871>.

151. Cangiarella J, Guth A, Axelrod D, et al. Is surgical excision necessary for the management of atypical lobular hyperplasia and lobular carcinoma in situ diagnosed on core needle biopsy?: a report of 38 cases and review of the literature. Arch Pathol Lab Med 2008;132:979-983. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18517282>.

152. Elsheikh TM, Silverman JF. Follow-up surgical excision is indicated when breast core needle biopsies show atypical lobular hyperplasia or lobular carcinoma in situ: a correlative study of 33 patients with review



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

of the literature. Am J Surg Pathol 2005;29:534-543. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15767810>.

153. Frouge C, Tristant H, Guinebreiere JM, et al. Mammographic lesions suggestive of radial scars: microscopic findings in 40 cases. Radiology 1995;195:623-625. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7753984>.

154. Margenthaler JA, Duke D, Monsees BS, et al. Correlation between core biopsy and excisional biopsy in breast high-risk lesions. Am J Surg 2006;192:534-537. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16978969>.

155. Parker SH, Burbank F, Jackman RJ, et al. Percutaneous large-core breast biopsy: a multi-institutional study. Radiology 1994;193:359-364. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7972743>.

156. Linell F, Pinder SE, Ellis IO. Precursor lesions of breast carcinoma. The Breast 1993;2:220-223. Available at: <http://www.sciencedirect.com/science/article/pii/096097769390003X>.

157. Middleton LP, Grant S, Stephens T, et al. Lobular carcinoma in situ diagnosed by core needle biopsy: when should it be excised? Mod Pathol 2003;16:120-129. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12591964>.

158. Sohn VY, Arthurs ZM, Kim FS, Brown TA. Lobular neoplasia: is surgical excision warranted? Am Surg 2008;74:172-177. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18306873>.

159. Bowman K, Munoz A, Mahvi DM, Breslin TM. Lobular neoplasia diagnosed at core biopsy does not mandate surgical excision. J Surg Res 2007;142:275-280. Available at: [http://www.journalofsurgicalresearch.com/article/S0022-4804\(07\)00187-4/abstract](http://www.journalofsurgicalresearch.com/article/S0022-4804(07)00187-4/abstract).

160. Kopans DB. LCIS found at core needle biopsy may not need surgical excision. AJR Am J Roentgenol 2008;191:W152; author reply W153. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18716082>.

161. O'Neil M, Madan R, Tawfik OW, et al. Lobular carcinoma in situ/atypical lobular hyperplasia on breast needle biopsies: does it warrant surgical excisional biopsy? A study of 27 cases. Ann Diagn Pathol 2010;14:251-255. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20637429>.

162. Rendi MH, Dintzis SM, Lehman CD, et al. Lobular in-situ neoplasia on breast core needle biopsy: imaging indication and pathologic extent can identify which patients require excisional biopsy. Ann Surg Oncol 2012;19:914-921. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21861212>.

163. Nagi CS, O'Donnell JE, Tismenetsky M, et al. Lobular neoplasia on core needle biopsy does not require excision. Cancer 2008;112:2152-2158. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18348299>.

164. Kaiser JS, Helvie MA, Blacklaw RL, Roubidoux MA. Palpable breast thickening: role of mammography and US in cancer detection. Radiology 2002;223:839-844. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12034957>.

165. Loving VA, DeMartini WB, Eby PR, et al. Targeted ultrasound in women younger than 30 years with focal breast signs or symptoms: outcomes analyses and management implications. AJR Am J Roentgenol 2010;195:1472-1477. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21098212>.

166. Robbins J, Jeffries D, Roubidoux M, Helvie M. Accuracy of diagnostic mammography and breast ultrasound during pregnancy and lactation. AJR Am J Roentgenol 2011;196:716-722. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21343518>.

167. Lehman CD, Lee CI, Loving VA, et al. Accuracy and value of breast ultrasound for primary imaging evaluation of symptomatic women 30-39 years of age. AJR Am J Roentgenol 2012;199:1169-1177. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23096195>.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

168. Lehman CD, Lee AY, Lee CI. Imaging management of palpable breast abnormalities. *AJR Am J Roentgenol* 2014;203:1142-1153. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25341156>.

169. Harvey JA, Mahoney MC, Newell MS, et al. ACR Appropriateness Criteria Palpable Breast Masses. *J Am Coll Radiol* 2016;13:e31-e42. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27814822>.

170. Soo MS, Rosen EL, Baker JA, et al. Negative predictive value of sonography with mammography in patients with palpable breast lesions. *AJR Am J Roentgenol* 2001;177:1167-1170. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11641195>.

171. Shetty MK, Shah YP. Prospective evaluation of the value of negative sonographic and mammographic findings in patients with palpable abnormalities of the breast. *J Ultrasound Med* 2002;21:1211-1216; quiz 1217-1219. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12418762>.

172. Moy L, Slanetz PJ, Moore R, et al. Specificity of mammography and US in the evaluation of a palpable abnormality: retrospective review. *Radiology* 2002;225:176-181. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12355002>.

173. Stavros AT, Thickman D, Rapp CL, et al. Solid breast nodules: use of sonography to distinguish between benign and malignant lesions. *Radiology* 1995;196:123-134. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7784555>.

174. Graf O, Helbich TH, Fuchsjäger MH, et al. Follow-up of palpable circumscribed noncalcified solid breast masses at mammography and US: can biopsy be averted? *Radiology* 2004;233:850-856. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15486217>.

175. Harvey JA, Nicholson BT, Lorusso AP, et al. Short-term follow-up of palpable breast lesions with benign imaging features: evaluation of 375 lesions in 320 women. *AJR Am J Roentgenol* 2009;193:1723-1730. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19933671>.

176. Shin JH, Han BK, Ko EY, et al. Probably benign breast masses diagnosed by sonography: is there a difference in the cancer rate according to palpability? *AJR Am J Roentgenol* 2009;192:W187-191. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19304679>.

177. Patterson SK, Neal CH, Jeffries DO, et al. Outcomes of solid palpable masses assessed as BI-RADS 3 or 4A: a retrospective review. *Breast Cancer Res Treat* 2014;147:311-316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25151294>.

178. Booi RC, Carson PL, O'Donnell M, et al. Characterization of cysts using differential correlation coefficient values from two dimensional breast elastography: preliminary study. *Ultrasound Med Biol* 2008;34:12-21. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2330278/pdf/nihms36784.pdf>.

179. Daly CP, Bailey JE, Klein KA, Helvie MA. Complicated breast cysts on sonography: is aspiration necessary to exclude malignancy? *Acad Radiol* 2008;15:610-617. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18423318>.

180. Huff JG. The sonographic findings and differing clinical implications of simple, complicated, and complex breast cysts. *J Natl Compr Canc Netw* 2009;7:1101-1104; quiz 1105. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19930976>.

181. Venta LA, Kim JP, Pelloso CE, Morrow M. Management of complex breast cysts. *AJR Am J Roentgenol* 1999;173:1331-1336. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10541113>.

182. Doshi DJ, March DE, Crisi GM, Coughlin BF. Complex cystic breast masses: diagnostic approach and imaging-pathologic correlation. *Radiographics* 2007;27 Suppl 1:53-64. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18180235>.

183. Hussain AN, Policarpio C, Vincent MT. Evaluating nipple discharge. *Obstet Gynecol Surv* 2006;61:278-283. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16551379>.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

184. Jain A, Crawford S, Larkin A, et al. Management of nipple discharge: technology chasing application. *Breast J* 2010;16:451-452. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20443787>.

185. Markopoulos C, Mantas D, Kouskos E, et al. Surgical management of nipple discharge. *Eur J Gynaecol Oncol* 2006;27:275-278. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16800258>.

186. Fuchsjager MH, Philipp MO, Loewe C, Helbich TH. [Diagnostic management of nipple discharge]. *Wien Klin Wochenschr* 2003;115 Suppl 2:33-39. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15518144>.

187. Jardines L. Management of nipple discharge. *Am Surg* 1996;62:119-122. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8554189>.

188. Ohuchi N, Furuta A, Mori S. Management of ductal carcinoma in situ with nipple discharge. Intraductal spreading of carcinoma is an unfavorable pathologic factor for breast-conserving surgery. *Cancer* 1994;74:1294-1302. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8055451>.

189. Gulay H, Bora S, Kilicturgay S, et al. Management of nipple discharge. *J Am Coll Surg* 1994;178:471-474. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8167884>.

190. Leis HP, Jr. Management of nipple discharge. *World J Surg* 1989;13:736-742. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2696228>.

191. Sabel MS, Helvie MA, Breslin T, et al. Is duct excision still necessary for all cases of suspicious nipple discharge? *Breast J* 2012;18:157-162. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22211878>.

192. Ashfaq A, Senior D, Pockaj BA, et al. Validation study of a modern treatment algorithm for nipple discharge. *Am J Surg* 2014;208:222-227. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24767970>.

193. Tokuda Y, Kuriyama K, Nakamoto A, et al. Evaluation of Suspicious Nipple Discharge by Magnetic Resonance Mammography Based on Breast Imaging Reporting and Data System Magnetic Resonance Imaging Descriptors. *Journal of Computer Assisted Tomography* 2009;33:58-62. Available at: http://journals.lww.com/jcat/Fulltext/2009/01000/Evaluation_of_Suspicious_Nipple_Discharge_by.10.aspx.

194. Ballesio L, Maggi C, Savelli S, et al. Role of breast magnetic resonance imaging (MRI) in patients with unilateral nipple discharge: preliminary study. *Radiol Med* 2008;113:249-264. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18386126>.

195. Foulkes RE, Heard G, Boyce T, et al. Duct excision is still necessary to rule out breast cancer in patients presenting with spontaneous bloodstained nipple discharge. *Int J Breast Cancer* 2011;2011:495315. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22295227>.

196. Dawood S, Cristofanilli M. What progress have we made in managing inflammatory breast cancer? *Oncology (Williston Park)* 2007;21:673-679. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17564325>.

197. Haagensen C. Inflammatory carcinoma. In: *Diseases of the breast*. 3rd ed. Philadelphia: WB Saunders; 1986. Available at:

198. Sakorafas GH, Blanchard DK, Sarr MG, Farley DR. Paget's disease of the breast: a clinical perspective. *Langenbecks Arch Surg* 2001;386:444-450. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11735019>.

199. Sakorafas GH, Blanchard K, Sarr MG, Farley DR. Paget's disease of the breast. *Cancer Treat Rev* 2001;27:9-18. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11237774>.



NCCN Guidelines Version 1.2017

Breast Cancer Screening and Diagnosis

200. Ikeda DM, Helvie MA, Frank TS, et al. Paget disease of the nipple: radiologic-pathologic correlation. Radiology 1993;189:89-94. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8396786>.

201. Smith RL, Pruthi S, Fitzpatrick LA. Evaluation and management of breast pain. Mayo Clin Proc 2004;79:353-372. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15008609>.

202. Noroozian M, Stein LF, Gaetke-Udager K, Helvie MA. Long-term clinical outcomes in women with breast pain in the absence of additional clinical findings: mammography remains indicated. Breast Cancer Res Treat 2015;149:417-424. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25556516>.

203. Barros AC, Mottola J, Ruiz CA, et al. Reassurance in the Treatment of Mastalgia. Breast J 1999;5:162-165. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11348279>.

204. Berg WA, Campassi CI, Ioffe OB. Cystic lesions of the breast: sonographic-pathologic correlation. Radiology 2003;227:183-191. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12668745>.

Discussion
Update in
progress